

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 November 2000 (30.11.2000)

PCT

(10) International Publication Number
WO 00/71701 A1

(51) International Patent Classification: C12N 15/10,
15/62, 15/64, C12P 21/02

(21) International Application Number: PCT/US00/14122

(22) International Filing Date: 23 May 2000 (23.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/135,677 24 May 1999 (24.05.1999) US

(71) Applicants (for all designated States except US): NEW
ENGLAND BIOLABS, INC. [US/US]; 32 Tozer Road,
Beverly, MA 01915 (US). BOSTON BIOMEDICAL RE-
SEARCH INSTITUTE [US/US]; 25 Grove Street, Water-
town, MA 02472 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): XU, Ming-Qun
[US/US]; 40 Crescent Road, Hamilton, MA 01982 (US).
EVANS, Thomas, C. [US/US]; 1st floor, 68 Albion Street,

Somerville, MA 02143 (US). PRADHAN, Sriharsa
[IN/US]; 6 Beverly Commons, Essex # 1, Beverly, MA
01915 (US). COMB, Donald, G. [US/US]; 9 Proctor
Street, Manchester, MA 01944 (US). PAULUS, Henry
[US/US]; 85 East India Row, Boston, MA 02210 (US).
SUN, Luo [CN/US]; 46 Plum Street, Hamilton, MA 01982
(US). CHEN, Lixin [CN/US]; 60 Rantoul Street #409,
Beverly, MA 01915 (US). GHOSH, Inca [IN/US]; 10
Cottage Avenue, Somerville, MA 02144 (US).

(74) Agent: WILLIAMS, Gregory, D.; New England Biolabs,
Inc., 32 Tozer Road, Beverly, MA 01915 (US).

(81) Designated States (national): AU, CA, CN, IN, JP, US.

(84) Designated States (regional): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR GENERATING SPLIT, NON-TRANSFERABLE GENES THAT ARE ABLE TO EXPRESS AN AC-
TIVE PROTEIN PRODUCT

(57) Abstract: A new type of transgene system is disclosed which allows efficient protein expression in a target host such as a
plant, but avoids the undesirable result of the migration of the transgene into related host systems and/or to the environment via the
pollen. The methods described herein may also be applied to the expression of virtually any protein of interest (e.g. a toxic protein)
in eukaryotic (yeast, insect, mammalian cells, etc.) and prokaryotic (*E. coli*, etc.) organisms.

BEST AVAILABLE COPY

**METHOD FOR GENERATING SPLIT, NON-TRANSFERABLE
GENES THAT ARE ABLE TO EXPRESS AN ACTIVE PROTEIN
PRODUCT**

5

BACKGROUND OF THE INVENTION

In the past few years, agriculture in the United States has been revolutionized by the introduction of transgenic crops that are resistant to specific diseases, insects, herbicides or have improved nutritional value. At the same time, much concern has been expressed around the world that these genetically modified (GM) agricultural products may be harmful to the consumer and that the transgenes could be transferred to related plant species so as to generate insect- or herbicide-resistant "superweeds" (Ferber, D., *Science* 286:1662 (1999)) or consumed by other organisms to their detriment (Losey, et al., *Nature* 399:214 (1999)). Whereas there is little scientific basis to the fear of harmful effects of "GM foods", the possibility that transgenes are transferred to other plants and thereby have an adverse ecological impact is not entirely unfounded (Bergelson, et al., *Nature* 395:25 (1998)). Such transfer could occur either by pollination of closely related species or by the transfer of gene fragments to unrelated plants by viral or plasmid vectors whose transmission may be mediated by plant-associated fungi, bacteria or insects.

There have been a number of techniques discussed for the prevention of transgene spread, however these

procedures either are designed to have a negative impact on the new hybrid plant (Gressel, *Trends Biotechnol.*, 17:361-366 (1999)), as in the case of tandem constructs or will not eliminate the possibility of spread by horizontal gene transfer (Bertolla and Simonet, *Res. Microbiol.*, 150:375-384 (1999)).

In this disclosure, we propose a new type of transgene that allows efficient protein expression but does not require a gene coupling approach and has a significantly lower chance of spread by horizontal gene transfer.

SUMMARY OF THE INVENTION

In accordance with the present invention, there is disclosed a new type of transgene system that allows efficient protein expression in a target host such as a plant, but avoids the undesirable result of the migration of the transgene into related host systems and/or to the environment via the pollen. The methods described herein can also be applied to the expression of virtually any protein of interest (e.g. a toxic protein) in eukaryotic (yeast, insect, mammalian cells, etc.) and prokaryotic (*E. coli*, etc.) organisms.

In each case, the target gene is split into at least two segments, each can be fused to a portion of an intein coding sequence. Each fusion gene is expressed as an inactive protein and these separately expressed fusion proteins are reassembled into an active form. Compartmentalization of

the gene fragments allows the target protein to be reconstituted in a desired location and can prevent the transmission of a functional gene to other organisms.

5 It should be noted that although the present invention is specifically exemplified in agriculture and plant biotechnology, the approach proposed here has a much broader scope and can be applied to any gene expressed in any organism for the prevention of its accidental transfer to
10 another organism.

DESCRIPTION OF THE DRAWINGS

Figure 1A - Protein Splicing Mechanism. Protein splicing
15 is a post-translational processing event involving the excision of an internal protein segment, the intein, from a precursor protein with the concomitant ligation of the flanking N- and C-terminal regions (the exteins). Sequence alignment reveals that there are highly conserved residues at the two splice
20 junctions: a cysteine or serine residue at the N-terminus of the intein, His-Asn at the C-terminus of the intein, and Cys, Ser or Thr as the first residue of the C-terminal extein. These conserved splice junction residues are directly involved in the catalysis of peptide bond cleavage and ligation of the protein
25 splicing reactions. The chemical mechanism of protein splicing with an intein which has cysteine residues at its N-terminus and adjacent to its C-terminus is shown in Figure 1: Step 1- Formation of a linear thioester intermediate by an N-S acyl

rearrangement of Cys1 at the N-terminus of the intein; Step
2- Formation of a branched intermediate by transesterification
involving attack by the Cys immediately following the C-
terminus of the intein on the thioester formed in Step 1; Step
3- Excision of the intein by peptide bond cleavage coupled to
succinimide formation involving the intein C-terminal Asn
residue; Step 4- Spontaneous S-N acyl rearrangement of the
transitory ligation product from a thioester to a stable amide
bond. Protein splicing involving other inteins presumably
proceeds by four analogous chemical steps, except that the
Cys residues shown in Figure 1 can be replaced by Ser or Thr,
so that Steps 1 and 4 are N-O and O-N acyl shifts,
respectively.

Figure 1B - Cartoon of protein splicing.

Figure 2 - Trans-Splicing.

Figure 2A-The association of the N-terminal and C-
terminal intein fragments aligns the two splice junctions for
the fusion of the N- and C-extein sequences. The splicing
reaction presumably occurs via the same splicing pathway as
the *cis*-splicing pathway proposed previously.

Figure 2B-Alternatively, in the absence of splicing the
intein could facilitate the association of the two extein
sequences with the subsequent generation of enzymatic
activity. This has been termed intein-mediated
complementation.

Figure 3 - *Ssp* DnaE intein gene arrangement in *Synechocystis* sp PCC6803. The genome of the blue-green algae *Synechocystis* sp PCC6803 contains the split *dnaE* gene with the fragments located 745 kb apart. The naturally occurring trans-splicing intein fuses the two gene product fragments to produce an active polymerase.

Figure 4A - Splitting of a target gene. A target gene can be split into two fragments with partial intein genes fused at the C- and N-terminal portions. These split genes can be placed into plant chromosomes so that the following expression can be reconstituted.

Figure 4B - Containment of a trans-gene. The gene of interest, in this case an herbicide resistance gene, is divided into two fragments (target N and target C) and an intein (IN_N and IN_C) is fused to each partial gene. The two gene fusions are placed on separate, remote locations on the genome. One of these may be in the chloroplast, the other in the nuclear genome. The chloroplast located transgene is transcribed and translated in the chloroplast while the nuclear transgene is transcribed in the nucleus and translated in the cytoplasm. Following translation of the nuclear gene it is transported into the chloroplast with the help of chloroplast transit peptide where it can associate with the other gene fragment using the intein as either an association or splicing element.

Figure 5 - Trans-splicing of acetolactate synthase (ALS) in *E. coli* strain ER2744. The target gene is split by intein fragments (IN_n and IN_c) and expressed as two inactive partial proteins. Protein trans-splicing produces an active target protein product in host cells.

Figure 6 - Sequence alignment for acetolactate synthase (ALS) genes (SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, SEQ ID NO:45 and SEQ ID NO:46). The gap region for *E. coli* acetolactate synthase II (ALSII) is underlined. The arrow indicates the split site for *E. coli* ALSII. The star indicates the split site for maize ALS.

Figure 7 - Plate assay showing that ALSIIIm-14 renders *E. coli* ER2744 resistant to valine and herbicide, SM. *E. coli* ER2744 cells were transformed with plasmid DNA expressing ALSII protein (1), ALSIIIm (2), ALSIIIm-14 (3) and plated on M9 medium containing 0.3 mM IPTG, with 100 µg/ml of valine (a), or with 100 µg/ml valine and 50 µg/ml SM (b). The plate assay was performed at 30°C for 50 hours.

Figure 8 - Production of recombinant ALSIIIm-14 through *Ssp* DnaE intein mediated *trans*-splicing. 2µl of whole cell extract, from cells transformed with expression plasmids for control (lane 1), ALSII (lane 2), ALSIIIm(N)-IN_n (lane 3), ALSIIIm(C)-IN_c (lane 4), ALSIIIm(N)-IN_n and ALSIIIm(C)-IN_c (lane 5), was run on an SDS-polyacrylamide (12%) gel, transferred

to a S&S nitrocellulose membrane, and probed with antiserum against ALSII N-terminus (Figure 8A) or against ALSII C-terminus (Figure 8B). (Figure 8C) The efficiency of *trans*-splicing is temperature sensitive. Western blot was performed using a antiserum against ALSII_m N-terminus. Protein extract was made from cells transformed with expression plasmids for control *E. coli* extracts contain a non-specific protein (the top band) that reacts with antiserum: (lane 1), ALSII (lane 2), ALSII_m(N)-IN_n and ALSII_m(C)-IN_c (lane 3 to lane 6). The cell culture temperature is 37°C for lane 1 to lane 3, 30°C for lane 4, 25°C for lane 5, and 15°C for lane 6.

Figure 9 - Assays for acetolactate synthase II (ALSII) Activity.

Figure 9A - Co-expression of ALSII_m(N)-IN_n and ALSII_m(C)-IN_c rescued cell growth on a valine plus herbicide added plate. *E. coli* ER2744, transformed with expression plasmids for ALSII (1), ALSII_m (2), ALSII_m(N)-IN_n and ALSII_m(C)-IN_c (3), ALSII_m(N)-IN_n (4), ALSII_m(C)-IN_c (5), ALSII_m(N) and ALSII_m(C) (6), were plated on M9 medium at 37°C (a), 37°C with 100 µg/ml valine (b), 30°C with 100 µg/ml valine (c), and 30°C with 100 µg/ml valine and 50 µg/ml sulfometuron methyl (SM) (d). Plates contained 0.3mM IPTG.

Figure 9B - Co-expression of ALSII_m(N)-IN_n and ALSII_m(C)-IN_c rescued cell growth in valine and herbicide added medium. *E. coli* ER2744, transformed with expression plasmids for fusion proteins as indicated under graph, was

cultured in M9 medium (0.3mM IPTG), with or without 100 µg/ml valine and 50 µg/ml sulfometuron methyl (SM) as indicated. OD₆₀₀ was taken to determine the cell growth rate after cells were cultured for 40 hours at 30°C.

5

Figure 9C - The time course study on the growth rate of cells expressing ALSII_m(N)-IN_n and ALSII_m(C)-IN_C. *E. coli* ER2744, transformed with the expression plasmids for proteins as indicated, was cultured at 30°C in M9 medium (0.3mM IPTG) with the addition of 100 µg/ml valine. The cell density was determined by measuring OD₆₀₀ at several time points as indicated.

10

Figure 10 - Western blot detection of *trans*-splicing product, maize ALS-14. 2 µl of whole cell lysate, from *E. coli* ER2744 cells transformed with expression plasmids for control (lane 1) (please note the antibody reacts with a non-specific protein in *E. coli*), cALS (lane 2), cALS(N)-IN_n (lane 3), cALS(C)IN_C (lane 4), cALS(N)-IN_n and cALS(C)-IN_C (lane 5), was run on a 12% SDS polyacrylamide gel, transferred to a S&S Nitracellulose membrane and probed with antiserum against cALS N-terminus (A) or cALS C-terminus (B). cALS indicates corn/maize ALS protein.

15

20

25

Figure 11 - Plating Assay for *Ssp* DnaE intein Cis-splicing Constructs. Plasmids pCE182DnaE, pCE215DnaE, pCE235DnaE, and pCE267DnaE encode for the 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) protein with the full

length *Ssp* DnaE intein inserted at amino acid positions 182, 215, 235 and 267, respectively. These were transformed into ER2799 *E. coli* cells (which require the EPSPS protein for viability in M9 minimal media), and plated on M9 minimal plates. Following incubation at 37°C overnight, individual clones on each plate were picked and stripped onto a single M9 minimal plate. This master plate was then incubated at 37°C overnight or RT for 2-3 days. As a control the pCYB3 plasmid was used as it carries no EPSPS gene, and there is no growth on the selection plate. pC+E2, a plasmid which contains the full length wild type EPSPS containing a Pro101Ser mutation, grows on M9 selection plate and also confers glyphosate resistance.

Figure 12 - Plating Assay for the *Ssp* DnaE intein Trans-splicing Constructs at Positions 215 and 235.

The activity of each 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) trans-splicing construct was assayed by co-transforming the matching constructs into *E. coli* ER2799 cells and plating on an M9 selection plate. pCYB3 or pKYB1 (New England Biolabs, Inc., Beverly, MA), which has no EPSPS gene present, was used to provide ampicillin or kanamycin resistance when testing the activity of each half of the EPSPS gene.

The plasmids used were: pC+E2, which contains the full length EPSPS mutant gene; p215EN2, which has the first 215

amino acids of EPSPS fused to the N-terminal splicing domain
of the *Ssp* DnaE intein; p235EN2, which has the first 235
amino acids of EPSPS fused to the N-terminal splicing domain
of the *Ssp* DnaE intein; pEPS#28, which contains amino acids
216-427 of the EPSPS gene fused to the C-terminal splicing
5 domain of the *Ssp* DnaE intein; pEPS#29, which contains
amino acids 236-427 of the EPSPS gene fused to the C-
terminal splicing domain of the *Ssp* DnaE intein; pEPS#33,
which has the first 235 amino acids of EPSPS fused to a
10 splicing defective N-terminal domain of the *Ssp* DnaE intein;
pEPS#37, which has amino acids 236-427 of EPSPS fused to
a splicing defective C-terminal domain of the *Ssp* DnaE intein;
pEPS#34, which has the first 235 amino acids of EPSPS, but
no intein fragment; and pEPS#36, which has amino acids 236-
15 427 of EPSPS and no intein fragment. These plasmids were
co-transformed, in various combinations, into ER2799 *E. coli*
cells, and plated on both LB plates and M9 plates, each plate
was supplemented with 100 µg/mL ampicillin and 50 µg/mL
kanamycin and 0.3 mM IPTG. Individual clones were picked
20 from each LB plate and stripped on one M9 selection plate
following incubation at 37°C overnight or RT for 2-3 days.
The M9 minimal media selection plate contained 100 µg/mL
ampicillin and 50 µg/mL kanamycin and 0.3 mM IPTG. The
combinations used were: WT, pC+E2 and pKYB; 215NC,
25 p215EN2 and pEPS#28; 215C, pEPS#28 and pCYB3; 235NC-
Dead, pEPS#33 and pEPS#37; 235NC, p235EN2 and
pEPS#29; 235N, p235EN2 and pKYB1; 235C, pEPS#29 and

pCYB3; 235N-215C, p235EN2 and pEPS#28; and 235 complement, pEPS#34 and pEPS#36.

Figure 13 - Glyphosate Resistance Liquid Assay for 235 Trans-splicing Constructs. The plasmid constructs were as described in Figure 12. The combinations used were: WT, pC+E2 and pKYB; 235NC-Dead, pEPS#33 and pEPS#37; 235NC, p235EN2 and pEPS#29; 235N, p235EN2 and pKYB1; 235C, pEPS#29 and pCYB3; and 235 complement, pEPS#34 and pEPS#36. These plasmids were co-transformed into ER2799 *E. coli* cells and plated on LB plates, supplemented with 100 µg/mL ampicillin and 50 µg/mL kanamycin; pCYB3/pKYB were co-transformed into *E. coli* ER2744, and plated on the LB plate, supplemented as described previously. A preculture was prepared for each transformation by inoculating the fresh colony into LB medium containing 100 µg/mL ampicillin and 50 µg/mL kanamycin at 30°C for overnight. Equal amounts of pre-culture (10-11 µL depending on the cell density) was inoculated into freshly-made M9 minimal medium containing 100 µg/ml of ampicillin, 50 µg/ml of kanamycin and 0.3 mM IPTG in the absence or presence of different amounts of glyphosate. The growth of each construct was measured by OD at 600 nm. Figure 13A, growth at 37°C. Figure 13B, growth at 30°C.

Figure 14 - Growth of the cis-splicing 235 construct in M9 liquid minimal media. A plasmid with the full length *Ssp DnaE* intein inserted into position 235 of 5-enolpyruvyl-3-

phosphoshikimate synthetase (EPSPS) was constructed. Two plasmid vectors were created (pCE235 DnaE and pEPS#31), one with a splicing competent *Ssp* DnaE intein (235 cis) and another with a splicing incompetent intein (235 dead). These plasmids were co-transformed with pKEB12 into ER2799 *E. coli* cells and plated on LB plates supplemented with 100 µg/mL ampicillin and 50 µg/mL kanamycin. A preculture was prepared for each transformation by inoculating the fresh colony into LB medium at 30°C for overnight. Equal amounts of pre-culture (10-11µL depending on the cell density) was inoculated into freshly-made M9 minimal medium containing 100 µg/ml of ampicillin, 50 µg/ml of kanamycin and 0.3 mM IPTG. The cell density was determined at various times using the OD at 600 nm.

Figure 15 is a table that shows the sites in the 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) protein that allow a 5 amino acid insertion and still result in active protein.

Figure 16 is a table that shows the sites in the 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) protein where a 5 amino acid insertion results in inactive protein.

Figure 17 is a map of pIH976. Circular double stranded DNA with a multiple cloning site. The restriction enzyme sites are indicated. Restriction sites with parenthesis are not

unique. Ptac represents tac promoter. Origin of replication is ori. This plasmid has tetracyclin drug resistant marker (Tetr).

Figure 18 is a map of pAGR3. Circular double stranded DNA (SEQ ID NO:76) with a multiple cloning site. The restriction enzyme sites are indicated below. Ptac represents Tac promoter. Origin of replication is ori. This plasmid has ampicillin drug resistant marker (amp^r). Lac operator and ribosome binding sites are indicated. Plasmid pAGR3 is an expression vector which includes several elements: (1) a synthetic tac promoter coupled to a symmetric synthetic lac operator sequence; (2) a lac ribosome binding site; (3) a polylinker for cloning with the ATG within the NcoI site being about seven nucleotides downstream of the ribosome binding site; (4) a copy of the lacI^q gene to provide repression of the tac promoter; (5) the replication origin from pBR322; (6) ampicillin resistance gene; and (7) a four-fold copy of the ribosomal transcription terminator upstream of the tac promoter. The transcription terminators lower the basal level of transcription by reducing read-through transcription from upstream promoters.

Figure 19 Trans-splicing of two unrelated gene products in *E. coli* using the *Ssp* DnaE intein as splice element.

Figure 19A Plasmid pIHaadE-N represents aadA gene (in black) fused to the N-terminal splicing domain of the *Ssp* DnaE intein (IN_n in grey). Plasmid pAGRE-CsmGFP plasmid represents the C-terminal splicing domain of the *Ssp* DnaE

intein (IN_C in grey) and smGFP (in black): The calculated molecular mass for each of the partners is indicated below in kDa. The arrow indicates a trans-splicing event resulting in a aadA-smGFP (57 kDa) fusion protein.

Figure 19B Ampicillin and spectinomycin sulphate selection of pIHaadE-N and pAGRE-CsmGFP plasmid in *E.coli* cells. *E.coli* were transformed with the plasmids indicated on the right side. Colony numbers are indicated on top.

Figure 19C Expression and detection of hybrid aadA-smGFP protein through trans-splicing. Western blot analysis of *E.coli* cell extracts expressing the constructs as indicated above the figure, using a monoclonal smGFP specific antibody. The relative positions of biotinylated MW markers (76, 57, 46, 37, 28 and 20) are in kDa. The protein bands corresponding to aadA-smGFP hybrid as well as IN_C-smGFP are indicated.

Figure 20 is a map of pNCT114/224. Circular double stranded DNA with a multiple cloning site capable of targeting gene/(s) to predetermined locus. The restriction enzyme sites are indicated. PpsbA and TpsbA represents photosynthetic polypeptide D1 gene promoter and terminator respectively. Origin of replication is ori. This plasmid has ampicillin drug resistant marker (amp^r). The homologous recombination sequences are indicated as left border (orf228-ssb for pNCT114 and 16SrDNA-trna^V for pNCT224) and right boarder

(orf1244 for pNCT114 and rps7/12 for pNCT224). CS represents the cloning sites.

Figure 21 Plant promoter PpsbA activity in *E.coli* and Trans-splicing of aadA and smGFP.

Figure 21A Plasmid p115ag/p225ag represents aadA gene (in black) fused to the *Ssp* DnaE intein N-terminal domain (IN_n in grey) and the *Ssp* DnaE intein C-terminal domain (IN_c in grey) fused to smGFP (in black). Both the hybrid genes are transcribed in opposite directions. The calculated molecular mass for each of the partner is indicated below in kDa. Arrow indicates a trans-splicing event resulting in a fused aadA-smGFP (57 kDa) protein.

Figure 21B Ampicillin and spectinomycin sulphate selection of p115ag and p225ag plasmid in *E.coli* cells. *E. coli* were transformed with the plasmids indicated on the right side. Colony identities are indicated on top. The digit after the plasmid is the isolate number. A plus symbol ("+") indicates the growth of the plasmid with the indicated antibiotics.

Figure 21C Expression and detection of hybrid aadA-smGFP protein through trans-splicing. Western blot analysis of *E. coli* cell extracts expressing the constructs as indicated above the figure, using a monoclonal smGFP specific antibody. The relative positions of biotinylated MW markers are to the

left in kDa. The protein bands corresponding to aad-smGFP hybrid as well as IN_C-smGFP are indicated.

Figure 22 Splicing *in cis* in plant cytoplasm. 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) and acetolactate synthase (ALS) genes are inserted in to the binary vector pBI121. The amino and carboxy terminal fragments of EPSPS or ALS are indicated in black. The *Ssp* DnaE intein (Intein) gene is flanked on either side by EPSPS/ALS fragment. Right and left boarder of the Agrobacterium is indicated as LB and RB. CaMV 35S promoter, NOS promoter (PNOS) and NOS terminator (TNOS) are indicated.

Figure 23 Nuclear transfer vector pBITPEC or pBITPECsmGFP. This binary vector has the CaMV35S promoter driving the rubisco3A transit peptide (TP) that is fused to the *Ssp* DnaE intein C-terminal splicing domain (IN_C). Genes to be cloned for organelle transport are indicated after IN_C. In case of pBITPECsmGFP the smGFP gene is cloned in to the multiple cloning site.

Figure 24 is the psbA promoter (PpsbA) sequence (SEQ ID NO: 59).

Figure 25 is the psbA terminator (TpsbA) (SEQ ID NO:60).

Figure 26 is the Rubisco3 transit peptide (SEQ ID NO:61). Nucleotides in lower case represent codon optimized units.

5 Figure 27 is the chloroplast gene targeting vector (pNCT114)(SEQ ID NO:62). Features of pNCT114 include: (1) vector backbone: pLITMUS28; (2) Inserted in *Bss*HII to *Bs*WI the left border, (orf228-ssb, 1210 bp) chloroplast genome targeting fragment; (3) inserted in *Avr*II to *Kpn*I the right border, (orf1244, 1550 bp) chloroplast genome targeting fragment; and (4) addition of *PpsbA* and *TpsbA* between *Bs*WI and *Pst*I, whereas the other pair is between *Avr*II and *Nco*I site.

15 Figure 28 is chloroplast gene targeting vector (pNCT224) (SEQ ID NO:63). Features of pNCT114 include: (1) vector backbone: pLITMUS28; (2) Inserted in *Bss*HII to *Bs*WI the left border, (16SrDNA-trnaV, 1680 bp) chloroplast genome targeting fragment; (3) inserted in *Avr*II to *Kpn*I the right border, (rps7/12, 1310 bp) chloroplast genome targeting fragment; and (4) addition of *PpsbA* and *TpsbA* between *Bs*WI and *Pst*I, whereas the other pair is between *Avr*II and *Nco*I site.

25 DETAILED DESCRIPTION OF THE INVENTION

Protein splicing involves the excision of an intervening sequence from a polypeptide with the concomitant joining of

the flanking sequences to yield a new polypeptide (Chong, et al., *J. Biol. Chem.*, 271:22159-22168 (1996)), as illustrated in Figure 1A and 1B. The elucidation of the mechanism of protein splicing has led to a number of intein-based applications (Comb, et al., U.S. Patent No. 5, 496,714; Comb, et al., U.S. Patent No. 5,834,247; Camarero and Muir, *J. Amer. Chem. Soc.*, 121:5597-5598 (1999); Chong, et al., *Gene*, 192:271-281 (1997), Chong, et al., *Nucleic Acids Res.*, 26:5109-5115 (1998); Chong, et al., *J. Biol. Chem.*, 273:10567-10577 (1998); Cotton, et al., *J. Am. Chem. Soc.*, 121:1100-1101 (1999); Evans, et al., *J. Biol. Chem.*, 274:18359-18363 (1999); Evans, et al., *J. Biol. Chem.*, 274:3923-3926 (1999); Evans, et al., *Protein Sci.*, 7:2256-2264 (1998); Evans, et al., *J. Biol. Chem.*, 275:9091-9094 (2000); Iwai and Pluckthun, *FEBS Lett.* 459:166-172 (1999); Mathys, et al., *Gene*, 231:1-13 (1999); Mills, et al., *Proc. Natl. Acad. Sci. USA* 95:3543-3548 (1998); Muir, et al., *Proc. Natl. Acad. Sci. USA* 95:6705-6710 (1998); Otomo, et al., *Biochemistry* 38:16040-16044 (1999); Otomo, et al., *J. Biolmol. NMR* 14:105-114 (1999); Scott, et al., *Proc. Natl. Acad. Sci. USA* 96:13638-13643 (1999); Severinov and Muir, *J. Biol. Chem.*, 273:16205-16209 (1998); Shingledecker, et al., *Gene*, 207:187-195 (1998); Southworth, et al., *EMBO J.* 17:918-926 (1998); Southworth, et al., *Biotechniques*, 27:110-120 (1999); Wood, et al., *Nat. Biotechnol.*, 17:889-892 (1999); Wu, et al., *Proc. Natl. Acad. Sci. USA* 95:9226-9231 (1998a); Wu, et al., *Biochim Biophys Acta* 1387:422-432 (1998b); Xu, et al., *Proc.*

Natl. Acad. Sci. USA 96:388-393 (1999); Yamazaki, et al., *J. Am. Chem. Soc.*, 120:5591-5592 (1998)).

Protein splicing *in trans* has recently been described both *in vivo* and *in vitro* (Shingledecker, et al., *Gene* 207:187 (1998), Southworth, et al., *EMBO J.* 17:918 (1998); Mills, et al., *Proc. Natl. Acad. Sci. USA*, 95:3543-3548 (1998); Lew, et al., *J. Biol. Chem.*, 273:15887-15890 (1998); Wu, et al., *Biochim. Biophys. Acta* 35732:1 (1998b), Yamazaki, et al., *J. Am. Chem. Soc.* 120:5591 (1998), Evans, et al., *J. Biol. Chem.* 275:9091 (2000); Otomo, et al., *Biochemistry* 38:16040-16044 (1999); Otomo, et al., *J. Biolmol. NMR* 14:105-114 (1999); Scott, et al., *Proc. Natl. Acad. Sci. USA* 96:13638-13643 (1999)) and provides the opportunity to express a protein as two inactive fragments that subsequently can undergo ligation to form a functional product (Figure 2).

Trans-protein splicing also occurs naturally in *Synechocystis* sp PCC6803 (Wu, H., et al., *Proc. Natl. Acad. Sci.* 95:9226 (1998)), where it is essential for forming a functional DNA polymerase III by joining two fragments of the DnaE protein, encoded by two genes separated by 750 kb of chromosomal DNA (Figure 3).

These observations led the present inventors to investigate whether a functional gene product could be generated by splitting the gene of interest into two fragments and fusing an intein fragment to each partial target gene.

Expression of the two protein fragments followed by *trans*-splicing, intein mediated complementation, or protein complementation would generate an active form of the target protein (Figure 4). In this scenario the target gene fragments can be located anywhere in the host genome, including being widely separated in the nucleus, chloroplast, mitochondria, plasmids, bacterial artificial chromosomes, yeast artificial chromosomes, or any combination of these. Furthermore, by placing the gene fragments into different organelles or plasmids, such as one half in the nucleus and the other half in the chloroplast or mitochondria of a plant, the transfer of both gene halves, needed to reconstitute the fully active target protein, for example, to a distant relative by pollination or by horizontal gene transfer via a bacterial, fungal, or viral vector would be virtually eliminated. This would greatly reduce and possibly eliminate the risk of the spread of a transgene outside of its relevant environment.

Two examples of splitting a target gene and reconstituting activity using a protein splicing element are described below. The two genes investigated were mutant forms of the acetolactate synthase (ALS) gene from *Escherichia coli* and the 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) gene from *Salmonella typhimurium*, which confer resistance to the sulfonylurea and glyphosate herbicides, respectively. Both enzymes are involved in the biosynthesis of protein building blocks. ALS is the first common enzyme in the biosynthesis of branched-chain amino

acids (LaRossa and Schloss, *J. Biol. Chem.*, 259:8753-8757 (1984); Chaleff and Ray, *Science*, 223:1148-1151 (1984); Falco and Dumas, *Genetics*, 109:21-35 (1985)) while EPSPS is required in the synthesis of aromatic amino acids (Stalker, et al., *J. Biol. Chem.* 260:4724-4728 (1985)). Inhibition of these enzymes by chemical compounds can lead to the death of the organism.

The commonly used sulfonylurea herbicides (SU), such as sulfometuron methyl (SM) (Short and Colburn, *Toxicol Ind. Health*, 15:240-275 (1999)), block the growth of bacteria, yeast and higher plants by inhibiting acetolactate synthase (ALS) (EC 4.1.3.18). In order to generate herbicide resistant plants, there was a great effort in identifying a mutant ALS gene which permits growth in the presence of SM. The mutations which render bacteria and yeast resistant to SM were the first to be reported (Hill, et al., *Biochem. J.*, 335:653-661 (1998)). Subsequently, similar point mutations were confirmed in the ALS genes isolated from naturally occurring resistant crops, corn, cocklebur and tobacco (Lee, et al., *EMBO J.*, 7:1241-1248 (1988); Bernasconi et al., *J. Biol. Chem.*, 270:17381-17385 (1995)). Some of these SU tolerant crops, such as corn ICI8532 IT and Pioneer 3180 IR have been commercialized.

In Example I below, the herbicide resistant gene was split and an intein fragment fused in-frame to each partial gene. The split gene was determined to confer resistance to

the herbicide SM in *E. coli*. *E. coli* was used as a model system since it contains the active ALSI and acetolactate synthase III (ALSIII) enzymes, but not an active ALSII. ALSI and ALSIII are the two isoforms of ALS genes in *E. coli* which are crucial for the synthesis of valine, isoleucine and leucine (DeFelice, et al., *Ann. Microbiol. (Paris)* 133A:251-256 (1982)). Their activity is sensitive to valine feedback inhibition. Therefore, by saturating the growth medium with valine, ALSI and III will be inhibited and the cells will stop growing. By introducing a recombinant ALSII into *E. coli* cells, their growth will be rescued since ALSII is resistant to valine inhibition. This feature makes *E. coli* strain ER2744 a good *in vivo* model system for investigating the activity of the *E. coli* ALSII gene genetically modified by a linker insertion or a trans-splicing intein element.

The second herbicide resistant gene tested was the *aroA* gene from *Salmonella typhimurium* that has a C301 to T mutation (Stalker, et al., *J. Biol. Chem.* 260:4724 (1985)). This encodes the 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) (EC 2.5.1.19) protein with a Pro101 to Ser change and is known to confer resistance to the herbicide glyphosate (commonly marketed as Round-Up®). In this embodiment, an N-terminal fragment of the EPSPS gene was fused to the N-terminal splicing domain of the *Ssp* DnaE intein and the C-terminal fragment of the EPSPS gene was fused to the C-terminal splicing domain of the *Ssp* DnaE intein. In order to determine the sites in the EPSPS protein that would tolerate

the insertion of an intein, a linker scanning experiment was performed (Biery, et al., *Nucleic Acids Res.*, 28:1067-1077 (2000)) (GPS®-LS from New England Biolabs, Inc., Beverly, MA) that randomly inserted 5 amino acids throughout the protein sequence. Inteins were inserted into those sites found to be tolerant of amino acid insertion. *Trans*-splicing constructs were then created that placed the gene fusion encoding the N-terminal fragment of EPSPS fused to the N-terminal domain of the *Ssp* DnaE intein on one plasmid and the C-terminal portion of EPSPS fused to the C-terminal splicing domain of the *Ssp* DnaE intein on another plasmid. For example the EPSPS protein could be split at the site corresponding to Gly235. The two plasmids were co-transformed into *E. coli* cells which lacked a functional EPSPS protein and cell growth on M9 minimal media in the presence or absence of the herbicide glyphosate was observed.

The activity of both the split ALS and the split EPSPS herbicide resistant genes were observed whether the intein was unmodified or had its catalytic residues changed, thus eliminating *trans*-splicing activity. This indicated that although splicing would generate a covalently attached protein product, it is not necessary to do so in every situation. The intein in this manifestation would work as an affinity domain to bring the two protein fragments together and in the correct orientation. In these experiments the presence of the intein was absolutely required for activity of the split proteins. This is based on the observation that both the split ALS and

EPSPS genes without an intein fusion were not able to allow *E. coli* growth on the appropriate herbicides.

In one embodiment of the invention, two gene fragments, fused to an intein splicing domain, are introduced independently into nuclear chromosomes, using selectable markers such as resistance to an antibiotic or other growth inhibitors to verify gene transfer. Independent transfer of the two fusion genes will assure a remote location on the plant genome, probably on separate chromosomes, thus excluding the possibility that both genes could be acquired by a single virus or plasmid vector for transfer to other organisms. If so desired, the remote location of the two genes can be assured by targeting to specific sites by homologous recombination with known DNA sequences.

In another embodiment, one of the two fusion proteins is transformed into the cell nucleus and the other into chloroplasts, so as to eliminate virtually any chance of gene transfer to related plants by any conceivable mechanism, including cross-pollination of related species, since only inactive fragments of the gene would be present in the pollen. The gene fragments in chloroplast are maternally transmitted and cannot be transmitted through pollen. The same consideration would apply to gene fragments expressed in mitochondria.

This technology may also be applied to non-plant systems. By way of example, a transgene to be compartmentalized could be split and an intein fused to the gene fragments. In the case of bacteria, the split genes are preferably placed far apart on the bacterial chromosome using standard chromosomal transformation techniques. As a further control measure the gene segments may also be arranged in opposite orientations. Another manifestation of this method is to split a target transgene in two and fuse to the appropriate intein domains prior to insertion of the split gene into a eukaryotic cell to prevent the transgene's activity from being spread to the environment or neighboring cells. The split gene is also placed far apart on the eucaryotic chromosome or placed on separate chromosomes. Furthermore, the gene fragments may be located in separate organelles such as the nucleus and mitochondria. The gene fragment in mitochondria is maternally transmitted.

One application of the present invention is in preventing the spread of complete transgenes to the environment from transgenic plants. This is accomplished by splitting the transgene fusions into two or more fragments and fusing these to intein fragments. The partial transgene fusions are located in separate compartments, such as one portion in the nuclear DNA and the second portion in the chloroplast DNA. Following expression of the partial genes, the protein fragments are directed to the site of activity where they associate to reconstitute the target protein activity. Only the

transgene fragment present in the nucleus is spread through pollen since the chloroplast DNA is passed to the next generation only maternally. This will vastly reduce the spread of the complete transgene to the environment.

5

Another advantage of the present invention is that the host cells expressing only one inactive fusion protein species of a protein can be handled safely, thereby reducing the risk of exposing humans and the environment to the target protein, which may be a toxin, etc. Also, splitting a target gene into two separate loci greatly reduces the chance of transferring the entire protein coding sequence into other organisms through DNA carriers (plasmid, virus, cosmid, etc.) or other means (cell fusion, etc.). One hypothetical case is to express a toxic gene, for example the diphtheria toxin. The diphtheria toxin protein is an extremely toxic protein to human and animal cells and needs to be handled extremely carefully. This protein has been tested in preclinical and clinical phase I trials for use as a drug to eradicate tumor cells (Kelley, *Proc. Natl. Acad. Sci. USA* 85(11):3980-3984 (1988); Alexander, *Neuron* 3(1):133-139 (1989); Maxwell, et al., *Cancer Res.* 51(16):4299-4304 (1991); Madshus, *J. Biol. Chem.*, 269(26):17723-17729 (1994); Murphy and vanderSpeck, *Semin Cancer Biol.* 6(5):259-267 (1995); Rozemuller and Rombouts, *Leukemia*, 12(5):710-717 (1998); Veggeberg, *Mol. Med. Today* 4(3):93 (1998); Kreitman, *Current Opin. Immunol.*, 11(5):570-578 (1999); Valleria, et al., *Protein Eng.* 12(9):779-785 (1999)). Therefore it would be advantageous to split the

10

15

20

25

diphtheria toxin gene into two intein fusion DNA segments and express them in two different bacteria or yeast strains. The two fusion proteins can be mixed, when it is needed, to assemble the toxin.

5

Thirdly, by compartmentalizing at least one of the fragments of the target gene into an organelle that is subject to maternal inheritance (e.g., chloroplasts or mitochondria), the genetic transfer of the functional gene to related organisms through processes such as cross-pollination can be avoided.

10

The invention described may also be utilized as a means for expressing any gene of interest in transgenic animals.

15

Transgenic animal models have been widely used as a scientific tool to conduct biomedical studies or to produce desired proteins. Transgenic mice and other transgenic animals, such as transgenic fish, frog, rat, cow, pig, etc. have been shown to express human genes (or a foreign gene) for research and commercial purposes, such as production of a vaccine or therapeutic agent, or used as an animal model for human disease (Alexander, *Neuron* 3(1):133-139 (1989); Groner, et al., *J. Physiol.* 84(1):53-77 (1990); Patil, et al., *Neuron* 4(3):437-447 (1990); Aloe, et al., *Growth Factors* 9(2):149-155 (1993); Aguzzi, et al., *Brain Pathol.* 4(1):3-20 (1994); Groner, et al., *Biomed. Pharmacother.* 48(5-6) 231-240 (1994); Schorderet, *Experientia* 51(2):99-105 (1995)). One of the concerns is that the transgenic animal may acquire an

20

25

undesired foreign gene and pass it on to the next generation and thereafter. This would result in genetically altered animal strains, which may have unforeseen social and ethical consequences. In accordance with the present invention, such a transgene can be split into two inactive fusion DNA fragments. One of them could be genetically integrated into an animal genome and the other fragment could be supplied by a DNA carrier (such as virus, etc.) which cannot be incorporated into the genome. Therefore, when one fusion protein from the animal and the other from the DNA carrier co-express, the fusion proteins will reassemble, trans-splice and produce an active protein. This gene arrangement can prevent animals from acquiring an intact foreign gene, thereby avoiding genetic contamination.

The compartmentalization of two gene fragments is an extension of trans-splicing. The protein in question is divided into fragments and the appropriate split genes separated onto the same or different DNA molecules. For example, the genes for the two halves of the DnaE protein from *Synechocystis* sp PCC 6803 with the *Ssp* DnaE or *Ssp* DnaB intein splicing domains fused to the appropriate fragments (Wu, et al., *Proc. Natl. Acad. Sci. USA*, 95:9226-9231 (1998a); Wu, et al., *Biochim Biophys Acta* 1387:422-432 (1998b)) could be divided so that one half is in the nucleus and the second half is in the mitochondria of a specified organism.

In carrying out the present invention, one must employ one or more of the following methods:

- (1) identifying a suitable split site on the target transgene;
- (2) the methodology for splitting the gene into two or more fragments and fusing each fragment to a split intein;
- (3) the methodology for successfully generating the split gene product into a functional enzyme or protein;
- (4) the methodology for screening the host cell for active gene product or organism;
- (5) location of split gene sequences in the relevant cellular compartment;
- (6) a method of splitting the target gene into more than two fragments;
- (7) use of protein complementation to present transgene spread; and
- (8) introduction of the transgene.

(1) A method for identification of a suitable split site on any transgene

One preferred method for identifying a split site on the transgene is based on the structural analysis of the protein of interest or its analogs and by sequence homology. This approach involves studying the known biochemical and X-ray, NMR or related structural information in order to determine a preferable intein insertion site and/or sites to divide the

protein into fragments. In particular, one should determine which are the pertinent reactive amino acid residues and their spacing and spatial arrangement within the protein. If possible, it may be ideal to split the target gene so that catalytic amino acids are distributed onto each fragment. This will increase the likelihood that neither fragment will have activity alone. The protein split site may be anywhere in the protein, but initial sites for testing should be loops or linkers present between secondary motifs such as beta sheets or alpha helices. The first loops chosen should not be part of the catalytic site, although the eventual split site may be located there. As a first trial, the preferred split site would be a loop or linker region between two folding domains within a protein. This increases the possibility that the protein fragments will fold properly when expressed separately.

If no biochemical or structural information is available for the protein of interest, then the alignment of similar protein sequences from different organisms or of similar protein sequences from the same organism may be informative. The protein alignment could be by sequence comparison by traditional methods or using any of a variety of computer programs such as GCG (Genetics Computer Groups, Madison, WI.) Regions of high conservation between similar proteins in all likelihood represent areas of general importance and splitting the protein in a region of high conservation should be reserved for later testing. Instead one should determine regions of low conservation, preferably regions that also vary

in amino acid number that lie between regions of high conservation. The low conservation indicates that there is a low probability of a catalytic residue being present and the variation in amino acid residue length indicates that the exact spacing between the conserved domains may not be dictated by this stretch of amino acids. These properties would be advantageous for a site of intein insertion and splitting a target protein.

Also, when choosing the site to insert an intein in the protein of interest one should test sites that possess amino acid residues favorable for the splicing activity of the intein being tested. Preferably a site in the target protein that was similar or identical to the naturally occurring extein residues of the intein under investigation could be chosen. Alternatively, residues known to facilitate proficient splicing may be inserted together with the intein. In this case, following the splicing reaction these residues would be present in the sequence of the spliced product and may alter the activity of the target protein. The effect of these extra residues on the target protein should be tested by inserting the extra amino acids into the target protein and checking for the desired property or activity.

Another preferred method is based on systematic scanning of a protein of interest by random linker insertion. Linker scanning can be performed by many methods (Gustin, et al. *Methods Mol. Biol.* 130:85-90 (2000); Hobson, et al.

Methods Mol. Biol. 57:279-285 (1996); Biery, *Nucleic Acids Res.* 28:1067-1077 (2000)). This protocol generates a library of genes with extra stretches of DNA randomly inserted throughout. When this library is translated it produces a set of proteins with extra amino acid residue(s) inserted in different positions. The library is then screened for the desired property of the target protein. For example, if the target protein confers resistance to an herbicide then the library is screened to determine which of the proteins with the extra amino acid residues can allow growth of the target organism in the presence of an herbicide. A list of sites in a protein that can tolerate extra amino acids is created. If structural or biochemical information is available, this list can be compared with the known information. An ideal case would involve choosing a split site that tolerates the extra amino acid insertion and is present in a linker or loop region and results in catalytic residues being located on different fragments. If no structural information is available then one would preferably begin by splitting the gene at the tolerant site closest to the middle of the target protein and continue testing split sites outward from there until the desired activity can be reconstituted. In both methods a preferred insertion site would also possess the native extein sequence for the intein being used, although this is not required. The fusion proteins may have optimized amino acid residues at the splice junctions that allow for a functional product to be assessed.

(2) A method for splitting a gene and fusing each gene fragment in-frame to a split intein coding sequence

5 Once a site to split a gene of interest has been determined (see above), then the target gene is split into two or more fragments using common genetic techniques (Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory, NY: Cold Spring Harbor Laboratory Press (1989)). For example, PCR primers, with appropriate restriction sites, may be designed so that one corresponded to the start of the target gene and the other to the sequence at the split site. Another set of PCR primers may be designed that correspond to the split site and the other end of the target gene. The two target gene fragments are then amplified by PCR (Sambrook, et al., *supra*) and cloned into a plasmid vector with the same unique cloning sites present in the PCR primers. Once cloned into separate vectors, intein fragments would be fused to the target genes.

10 In one method, the C-terminal end of DNA coding for an N-terminal portion of the target protein would be fused to the N-terminal end of the DNA coding for an N-terminal portion of the intein, and -in a separate fusion- the N-terminal end of DNA coding for a C-terminal portion of the target protein

15 would be fused to the C-terminal end of DNA coding for a C-terminal portion of the intein.

20 These gene fragment fusions are then transferred to the same or separate expression vectors and transformed

into bacterial or eucaryotic cells, existing as single or multicellular organisms, to screen for the desired activity of the target protein. It should be noted that the gene fragments in question could be cloned using restriction sites within or external to the intein gene present either naturally or added by mutation. Also, recombination sites may be used instead of restriction enzyme sites for the movement of the gene by recombination. The gene or gene fragments may then be transferred and/or expressed from a plasmid vector, a viral genome or the genome of a bacterial, eucaryotic, or archeal organism. One preferred method is to utilize a naturally occurring *trans*-splicing intein, for example the intein from the *dnaE* gene of *Synechocystis* species PCC6803 (Wu, et al., *Proc. Natl. Acad. Sci. USA* 95:9226-9231 (1998)). However, any of the known inteins could be used (See InBase at http://www.neb.com/neb/frame_tech.html; Perler, et al., *Nucleic Acids Res.*, 28:344-345 (2000)). This would involve splitting the full length intein in order to generate the desired affinity or *trans*-splicing domains. One method would be to split the full length intein in the linker region between the blocks B and F of the protein splicing domains (Petrokovski, *Protein Sci.* 7:64-71 (1998); Perler, et al., *Nucleic Acids Res.* 25:1087-1093 (1997); Perler, et al., *Nucleic Acids Res.*, 28:344-345 (2000)).

(3) Creating a functional protein from expressed split fragments

The next step is to use an intein as an affinity domain to facilitate complementation and reconstitution of the N- and C-terminal halves of a protein into a functional enzyme. The sites to determine protein splitting would be as described in (1) above and the cloning of the target gene fragments and the addition of the intein domains as described in (2). In this case the intein fragments need not cause splicing of the two protein fragments to reconstitute enzyme activity. In one preferred embodiment, the intein domains would be mutated to abolish the possibility of splicing activity and would act only as a facilitator of protein complementation. The intein splicing activity could be abolished by mutating the amino acid residues involved in the splicing reaction (Xu, et al., *EMBO J.* 15:5146-5153 (1996); Chong, et al., *J. Biol. Chem.* 271:22159-22168 (1996); Chong, et al., *Biochem. Biophys Res. Commun.*, 259:136-140 (1999); Chong, et al., *Gene*, 192:271-281 (1997); Chong, et al., *Nucleic Acids Res.*, 26:5109-5115 (1998); Chong, et al., *J. Biol. Chem.*, 273:10567-10577 (1998); Chong and Xu, *J. Biol. Chem.*, 272:15587-15590 (1997); Evans, et al., *J. Biol. Chem.*, 274:18359-18363 (1999); Evans, et al., *J. Biol. Chem.*, 274:3923-3926 (1999), Evans, et al., *Protein Sci.*, 7:2256-2264 (1998); Evans, et al., *J. Biol. Chem.*, 275:9091-9094 (2000); Mathys, et al., *Gene*, 231:1-13 (1999); Paulus, *Chem. Soc. Rev.*, 27:375-386 (1998); Perler, et al., *Nucleic Acids Res.*, 25:1087-1093 (1997); Pietrokovski, et al., *Protein Sci.*, 3:2340-2350 (1994); Pietrokovski, et al.,

Protein Sci., 7:64-71 (1998), Scott, *Proc. Natl. Acad. Sci. USA*, 96:13638-13648 (1999), Shingledecker, et al., *Arch Biochem. Biophys.* 375:138-144 (2000); Southworth, et al., *Biotechniques* 27:110-120 (1999); Telenti, et al., *J. Bacteriol.*, 179:6378-6382 (1997); Wood, et al., *Nat. Biotechnol.*, 17:889-892 (1999); Wu, et al., *Biochim Biophys Acta* 1387:422-432 (1998b); Wu, et al., *Proc. Natl. Acad. Sci. USA* 95:9226-9231 (1998a)).

In another embodiment the intein affinity domain could retain its normal catalytic residues. Furthermore, the intein may be comprised of a deletion or mutant form such that it is significantly smaller or larger or contains non-native amino acid residues when compared to its original primary sequence. The deletion forms of the intein could be created by sequentially decreasing the size of the intein either at the gene level or proteolytically and then testing for affinity activity. The affinity activity could be tested by using the split herbicide resistant gene and fusing the new deletion mutant to the appropriate herbicide resistant gene fragments and looking for growth on the herbicide in question. Mutants of the intein fragment could be formed by error prone PCR, linker scanning, site directed mutagenesis, or by mutagenic compounds and the activity of the intein fragments tested as described above. Note the herbicide resistance gene could be substituted by a drug resistance gene, green fluorescent protein or any selectable marker. The affinity of the intein fragments could also be tested by immobilizing one fragment

on a solid support and testing for the binding of the second fragment to the first fragment.

(4) A method of screening for constructs producing active proteins of interest in a suitable host cell or organism

The screen for the target gene activity will vary with the target gene but could be by *in vitro* assay following expression and purification or in a crude cell lysate or *in vivo* by determining protein activity by cell phenotype, such as viability, morphology, sensitivity, or insensitivity to a drug or compound, appearance, or ability to bind or not bind a specific molecule or compound. One preferred method is to use *E. coli* as host cells to test, for example, herbicide resistant activity of the re-assembled product of a split gene. The *E. coli* cells must be sensitive to the herbicide in question. The target gene fragments, with the intein fusion, is present on a plasmid or plasmids and is transformed into *E. coli* cells using standard techniques.

The gene fusions are expressed either constitutively or by an inducible promoter. *E. coli* are then tested for growth under selection conditions, i.e. in the presence of herbicide, in both the presence or absence of the appropriate gene fragments. Growth in the presence of the gene fragments indicates the reconstitution of the target protein activity. The *E. coli* cells could be substituted with any bacterial, archaea,

or eucaryotic cell type (either single or multicellular) as well as a virus by employing techniques well known in the art.

Furthermore, both of the target gene fragments could
5 be present in the genome of the organism, or one fragment could be present in the genome and the other in a plasmid or some other vector. The target protein fragments could be expressed in one organism together or separately and added to another cell type for assay. The fusion could be tested
10 directly in plant cells or other multicellular organisms by placing the transgene fragments in the host organisms nuclear, chloroplast, or mitochondrial genome and determining if the desired activity is present. The target gene or protein fragments could be delivered by a bacterial, fungal,
15 viral, micellar, mechanical (biolistic) or similar vector to the cell type or organism to be tested.

(5) Location of split genes

20 The present invention also comprises location of the split target gene sequences in different cellular compartments, different locations on the chromosome, or different vectors. One preferred method is to position the two split gene sequences in the nucleus, chloroplast, mitochondria, bacterial
25 artificial chromosome, yeast artificial chromosome, plasmid, preferably not both in any one of the aforementioned. Location of fragments can be accomplished in accordance with standard molecular biology techniques. In order to

reconstitute the gene product from its fragments, the appropriate gene fragments must be fused to a targeting/localization sequence so that their protein products are transported into a cellular compartment (e.g., the chloroplasts) where functional reconstitution can occur.

(6) A method of splitting the target gene into two or more fragments

The present invention also embodies methods for splitting the target gene into two or more fragments and reconstituting the desired activity by trans-splicing, intein mediated complementation or protein complementation of all the necessary fragments. For example, inteins with differing affinities could be attached to the target protein fragments so that they reassemble the active protein, in a manner described previously (Otomo, et al., *Biochemistry*, 38:16040-16044 (1999); Otomo, et al., *J. Biomol. NMR*, 14:105-114 (1999)). In this case each fragment could be located far apart in the chromosome, on a separate chromosome or in multiple locations as described above, except that the number of locations could match the number of fragments the protein was divided into.

(7) Use protein complementation in the prevention of transgene spread

This protocol uses the natural complementation activity of two protein fragments to reconstitute the desired protein property. The two genes encoding the protein halves may be

located in the nucleus, chloroplast, mitochondria, bacterial
artificial chromosome, yeast artificial chromosome, plasmid or
any combination of those organelles or vectors. Following
expression, both protein fragments may be targeted to the
5 site of protein action and the desired protein property
generated by complementation of the protein fragments.
Protein complementation has been reported previously (Rossi,
et al., *Trends Cell Biol.* 10:119-122 (2000)) and so makes a
viable alternative to using an intein as a complementation
10 domain. The procedures necessary to carry out this
experiment are similar to what has already been discussed
except no intein fusion is used. A site to split a target gene is
determined as described in (1). The transgene fragments are
cloned as described in (2), except that an intein is not used
15 as a fusion partner. The screening for activity of the split
protein is conducted as described in (4).

(8) Introducing a transgene into an organism by viral infection

20 In yet another embodiment, the two transgene
fragments, either intein fusions or not, may be packaged into
separate viral particles. These viruses co-infect an organism
and both transgenes are expressed. The desired protein
25 property is generated following protein splicing, intein
mediated complementation, or protein complementation. One
preferred method comprises choosing the split site, clone the
fragments and check for activity *in trans* as described in (1),
(2), and (4). The appropriately split transgene or transgene-

intein fusions are packaged into adenovirus. The adenoviruses containing the appropriate transgenes can be introduced into a subject organism and upon transfection introduce the two gene fragments so that the target protein activity can be expressed.

BRIEF DESCRIPTION OF THE EXAMPLES

In Example I, we demonstrate a method of splitting a herbicide resistant gene by an intein. We show how to select potential split sites in the *E. coli* herbicide resistant gene encoding for acetolactate synthase (ALS) based on the sequence homology analysis and the crystal structure of the protein of interest or its analog. The DNA fragment encoding for the N-terminal 327 amino acid residues of the ALS protein was fused in frame to the N-terminal 123 amino acids of the *Ssp* DnaE intein while the DNA fragment encoding for the C-terminal 221 amino acid residues was fused in frame to the C-terminal 36 amino acids of the *Ssp* DnaE intein. A plasmid vector bearing one of the fusion genes was expressed as an inactive ALS protein fragment. When both fusion gene vectors were introduced into the same host cell and co-expressed, the two inactive fusion proteins underwent trans-splicing to produce a functional enzyme *in vivo*, conferring herbicide resistance to the *E. coli* host cells. This approach may be applied to selection of suitable sites in any gene for fusion to an intein sequence.

In Example II, we demonstrate how to choose a split site in the maize ALS gene based on the sequence homology of the maize ALS gene and its *E. coli* counterpart, ALSII gene. The DNA encoding the N-terminal 397 amino acid residues of the maize ALS gene was fused in-frame to the DNA sequence encoding the N-terminal 123 amino acids of the *Ssp* DnaE intein while the DNA fragment encoding the C-terminal 241 amino acid residues was fused in frame to the DNA encoding the C-terminal 36 amino acids of the *Ssp* DnaE intein. We show that, when the two fusion genes were co-expressed, the two fusion proteins underwent trans-splicing to produce a protein product of expected size for the mature protein.

In Example III, we demonstrate a method of identifying potential split sites in a mutant *S. typhimurium aroA* gene encoding 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) based on transposon random linker insertion. Two sites at amino acid positions 215 and 235 of EPSPS among all 42 potential sites were chosen to split the EPSPS gene. The DNA fragment encoding the N-terminal 215 or 235 amino acid residues of the EPSPS protein was fused in-frame to the N-terminal 123 amino acids of the *Ssp* DnaE intein while the DNA fragment encoding the C-terminal 212 or 192 amino acid residues of EPSPS was fused in-frame to the DNA encoding the C-terminal 36 amino acids of the *Ssp* DnaE intein. When only introducing half of the EPSPS gene with or without the intein fused and the two complement halves without intein

into ER2799, the EPSPS was expressed as a non-functional protein. However, when introducing both the halves of EPSPS fused with both active or inactive intein halves into ER2799, the EPSPS was expressed as a functional protein and confers resistance to the herbicide glyphosate indicating that the N- and C-terminal halves of the *Ssp* DnaE intein facilitate the complementation and reconstitution of the N- and C-terminal halves of the EPSPS protein by bringing the EPSPS halves in close proximity.

In Example IV, we describe a method in which two unrelated gene products such as aminoglycoside-3-acetyltransferase (enzyme responsible for metabolism of drug spectinomycin or streptomycin) and *Aequorea victoria* soluble modified green fluorescent protein could be trans-spliced to one hybrid protein in *E.coli* cell. Both the genes are located on two different plasmids with respective trans-splicing elements from *Ssp* DnaE intein. The plasmids have two independent mechanisms of expression. This hybrid protein confers resistance to spectinomycin sulphate.

In Example V, we describe a method in which two unrelated genes, such as *aadA* (encodes for aminoglycoside-3-acetyltransferase) and smGFP (soluble modified green fluorescent protein), could be located on a single *E.coli*-plant binary vector under the transcriptional and translational control by a chloroplast promoter (*PpsbA*). Both the genes when expressed are capable of producing a hybrid

aminoglycoside-3-acetyltransferase- soluble modified green fluorescent protein. Thus this method allows for rapid trans-splicing screening of protein/protein fragments before introducing to the plant cells using promoter that could be recognized both by *E.coli* and plant cellular machinery.

In Example VI, we describe a method in which a cis-splicing construct containing two fragments of either 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) or acetolactate synthase (ALS) genes along with a *Ssp* DnaE intein is capable of splicing into a mature protein in plant cytoplasm. This experiment will enforce the idea of cis/trans-splicing in the cytoplasm. This technique would be useful for proteins, which need specific modification for activity/folding in cytoplasmic environment. A part of the target protein gene with necessary transport signal and splicing element will be placed in an organelle for cytoplasmic transport in the form of a pre-cursor polypeptide.

In Example VII, Section 1, we describe a method in which two unrelated genes, such as *aadA* (encodes for aminoglycoside-3-acetyltransferase) and *smGFP* (soluble modified green fluorescent protein), could be located on the chloroplast genome and produce a hybrid protein via protein trans-splicing. Success in this method will lead to compartmentalization of protein/protein fragments and trans-splicing of the functional protein. Also transformation of several separated genes in one vector to form a

multifunctional protein simplifying engineering of novel characters.

5 In Example VII, Section 2, we describe a method in which two unrelated genes/gene fragments could be localized in two different compartments in plant cell, such as chloroplast and nucleus and express the respective protein/polypeptide. The nuclear encoded component is tripartite with a chloroplast transit peptide which will help the
10 protein fragment to be synthesized in cytoplasm and migrate in to the chloroplast for the trans-splicing event to occur. The chloroplast half will be as an integrated component in the circular genome of the organelle. The resulting plants will not be able to transfer the novel character of the newly
15 introduced transgene to any closely related species.

The present invention is further illustrated by the following Examples. These Examples are provided to aid in the understanding of the present invention and are not
20 construed as a limitation thereof.

The references cited above and below are hereby incorporated by reference.

EXAMPLE I

Production of Functional Herbicide-resistant Acetolactate Synthase in *E. coli* by Protein *Trans*-splicing

5 In this Example we demonstrate a method to split the gene which encodes *E. coli* acetolactate synthase II (ALSII; EC 4.1.3.18; acetohydroxyacid synthase), possessing a herbicide-resistant mutation (Yadav *et al*, *Proc. Natl. Acad. Sci.*
10 *USA*, 83:4418-4422 (1986); Hill *et al.*, *Biochem. J.*, 335:653-661 (1998)), by fusion with *Ssp* DnaE intein coding sequences (Evans *et al*, *J. Biol. Chem.* 275:9091-9094 (2000); Scott, *et al.*, *pro. Natl. Acad. Sci. USA*, 96:13638-13643 (1999)). We were able to reconstitute a functionally active ALSII enzyme
15 through protein *trans*-splicing in the bacterium *E. coli* ER2744 (fhuA2 glnV44 el4- rfbD1? relA1? endA1 spoT1? thi-1 Δ (mcrC-mrr)114::IS10 lacZ::T7 gene1) (Figure 5). First, we show how to select a potential split site in the acetolactate synthase II gene based on the analysis of its sequence and structure
20 homology. Then we show how to design and carry out experiments to analyze the protein *trans*-splicing activity of the split ALS protein and how to assay the enzymatic activity of reconstituted ALS. We demonstrate that the two portions of the ALS fusion protein, produced from two separate
25 plasmid vectors, undergo *trans*-splicing to produce a protein product of expected size for the mature protein. Furthermore, co-expression of the split ALS gene fragments conferred resistance to a herbicide in the *E. coli* ER2744. This method

may be applied to the production of any protein of interest utilizing *trans*-splicing inteins.

1. Cloning of wild-type *E. coli* ALSII and generation of its herbicide resistant mutant

The initial step is to clone the wild type ALSII and to create a herbicide resistant ALSII mutant carrying Alanine26 to Valine substitution (Yadav *et al*, *Proc. Natl. Acad. Sci. USA*, 83:4418-4422 (1986); Hill *et al*, *Biochem. J.*, 335:653-661 (1998)). *E. coli* strain MI162, containing an enzymatic active copy ALSII, was obtained from CGSC, *E. coli* Genetic Stock Center (Yale University, New Haven, CT). Genomic DNA was extracted from *E. coli* strain MI162 using QIAamp Tissue Kit (Qiagen, Inc., Studio City, CA). DNA Polymerase Chain Reaction (PCR) was performed on the *E. coli* DNA sample to clone the full length ALSII using primers 5'-GGACGGGGAAGTAACTATG-3' (SEQ ID NO:1) and 5'-CCACGATGACGCACCACGCG-3' (SEQ ID NO:2) and Vent® DNA Polymerase (New England Biolabs, Beverly, MA). The ALSII coding sequence was further amplified using primers 5' GGAGGGGGCATATGAATGGCGCACAGTGGG-3' (SEQ ID NO:3) and 5'-GGGGGGTCATGATAATTTCTCCAAC-3' (SEQ ID NO:4) and cloned into *Nde*I and *Pst*I sites of pTYB1 plasmid (New England Biolabs, Beverly, MA), creating a vector, pALSII. A shorter construct, pTYBT-ALSII, was obtained by the removal of a 3-kb non-essential sequence from pALSII by restriction digestion with *Pme*I and BstZ172 followed by self ligation. The herbicide resistant mutation, Alanine26 to Valine,

was introduced in pTYBT-ALSII by site-directed mutagenesis using Quickchange Site-Directed Mutagenesis kit (Stratagene, La Jolla, CA). The mutagenesis primers were 5'-CCGGGTGGCG TAATTATGCCGGTTTACG-3' (SEQ ID NO:5) and 5'-CGTAAACCG GCATAATTACGCCACCCGG-3' (SEQ ID NO:6). The mutated ALSII (ALSII_m) coding sequence generated by partial *Nde*I and *Pst*I digestion of pTYBT-ALSII_m was ligated with pTYB1 to produce an ALSII_m expression vector, pALSII_m.

2. Selection of Split Site

One preferred method for identifying a suitable split site within any gene, is to analyze the sequence homology of a family of proteins and to examine its protein structure or the structure of its homologues (Ibdah *et al.*, *Biochemistry*, 35:16282-16291 (1996)). Sequence alignment and structure comparison suggest that the ALS genes of bacteria, yeast and higher plants share highly conserved regions (Figure 6, only partial sequence alignment is shown here). Still, there are highly variable regions present in the proteins, such as the region around amino acid residues Q327 and C328 in the isoform II of the *E. coli* acetolactate synthase (Figure 6). *E. coli* ALSII has a 10 amino acid gap in this region compared to other homologues and the flanking sequence has less homology among ALS genes from different species (Figure 6). Furthermore, analysis of the crystal structure of a homologue, pyruvate oxidase, suggests that Q327 and C328 are likely to be located in a linker structure between two intra-molecular

domains, away from the catalytic core (Ibdah *et al.*,
Biochemistry, 35:16282-16291 (1996)). We reasoned,
therefore, that ALSII split by an intein at this region may
retain the necessary flexibility to allow efficient protein *trans*-
splicing. In addition, insertion of a foreign protein sequence
into this location may have less or no effect on the structure
of the catalytic domain of ALSII and its enzymatic activity.
Thus amino acid residues Q327 and C328 were selected as
one of the split sites for *E. coli* ALSII (indicated by an arrow,
Figure 6).

3. *E. coli* assay system

The isoform II of the *E. coli* acetolactate synthase that
possesses the mutation Ala26Val, referred to as ALSII_m,
confers resistance to sulfonylurea herbicides (SU), such as
sulfometuron methyl (SM), in *E. coli* strain ER2744. *E. coli*
ER2744 strain was employed as an *in vivo* model system for
assessing the activity of the herbicide resistant *E. coli* ALSII
gene, genetically modified by a linker insertion between Q327
and C328. *E. coli* ER2744 is derived from wild type *E. coli* K12
that contains the active ALSI and ALSIII enzymes, but not an
active ALSII. ALSI and ALSIII are two isoforms of ALS genes in
E. coli, which are crucial for the synthesis of valine, isoleucine
and leucine (LaRossa and Schloss, *J. Biol. Chem.* 259:8753-
8757 (1984)). Their activity is sensitive to the valine feedback
inhibition. Therefore, by saturating the growth medium with
100 µg/ml valine (Sigma, St. Louis, MO), ALSI and III will be

inhibited and the cells will stop growing. By introducing a recombinant herbicide resistant ALSII (ALSII_m) into *E. coli* cells, their growth will be rescued since ALSII is resistant to valine inhibition.

4. Generation of a modified herbicide resistance ALS gene

Inteins often require certain amino acid residues flanking its N- and C-termini to achieve optimal splicing or *trans*-splicing activity. For example, the intein from the *dnaE* gene of *Synechocystis* species PCC6803 spliced efficiently when 5 native residues were present at both its N- and C-termini, while deletion of these residues inhibited splicing activity to various extents (Evans *et al.*, *J. Biol. Chem.* 275:9091-9094 (2000)). Inclusion of these optimal amino acid residues at the splice junctions may be required for proficient splicing activity. The resulting product may therefore possess these residues at the ligation junction of two protein sequences. Thus, for each intein insertion site, it is necessary to assess if these extra amino acid residues will have an adverse effect on the activity of the product.

ALSII_m-14 was constructed by insertion of a synthetic DNA linker (New England Biolabs, Beverly, MA), encoding the following 14 amino acid residues (NH₂-LEKFAEYCFNKSTG-COOH (SEQ ID NO:7)), into the ALSII_m coding sequence between Q327 and C328A. The herbicide resistance activity of ALSII_m-14 was examined using *E. coli* ER2744 host cells

transformed by the plasmid expressing ALSII_m-14 protein. *E. coli* ER2744 cells transformed with plasmids expressing the wild type ALSII and herbicide resistant ALSII (ALSII_m) were used as controls.

5 Plate assays were conducted to examine the capability of ALSII_m-14 to rescue *E. coli* ER2744 from valine (100 µg/ml) or valine plus herbicide SM (50 µg/ml, Supelco Park, Bellefonte, PA) saturated M9 minimum medium plate
10 (Sambrook *et al.*, (1989)). The M9 medium contains 2 µg/ml Thiamin, 2 mM MgSO₄, 0.1 mM CaCl₂, 0.2% glucose, 50 µg/ml of kanamycin, 100 µg/ml of ampicillin and 0.3 mM IPTG. For the plating assay, 100 µl of 25 mg/ml Valine with or without 50 µl of 25 µg/ml Sulfometuron methyl (SM) was spread on M9
15 selection plate. To assay bacterial growth, overnight cultures were streaked on M9 plates with or without valine and/or SM. The plates were incubated at various temperatures (as indicated in Figure 7) for 48 to 72 hrs before the pictures were taken. On the plate supplemented with valine, cells
20 expressing either ALSII, ALSII_m or ALSII_m-14 were able to grow (Figure 7-a). However, when both valine and SM was applied, only strains expressing herbicide-resistant ALSII_m or ALSII_m-14 were able to grow (Figure 7-b). These *in vivo* results demonstrated that ALSII_m with 14 amino acid
25 residues inserted at the proposed split site, rescued *E. coli* ER2744 growth in the presence of valine and SM. Therefore, ALSII_m-14 is functionally active and the 14 amino acid insertion does not affect its enzymatic activity.

5. Construction of ALSII-Intein fusion genes

Next, the *E. coli* ALSII_m gene was split and fused in-frame to the N- and C-terminal halves of the *Ssp* DnaE intein coding regions. The fusion genes were created using two compatible *E. coli* expression vectors, pMEB10 and pKEB1, which are capable of co-expressing two intein fusion genes in the same *E. coli* host cell, as previously described by Evans et al. (*J. Biol. Chem.* 275:9091-9094 (2000)). The DNA sequence encoding for an N-terminal fragment of 327 amino acids of the herbicide resistant ALSII (ALSII_m) gene was fused in frame to the coding region for the 7 amino acid residues flanking the N-terminus of the *Ssp* DnaE intein, followed by the intein N-terminal 123 amino acid residues (IN_n) (Figure 5). The DNA sequence encoding the C-terminal 221 amino acid residues of ALSII_m was fused in frame to the DNA sequence encoding the C-terminal 36 amino acid residues of the *Ssp* DnaE intein (IN_c) and the 7 amino acid residues flanking the C-terminus of the intein (Figure 5). ALSII N-terminal fragment was amplified from pALSII_m using primers 5'-GGGGGTCATGAATGGCGCACAGTGGG-3' (SEQ ID NO:10) and 5'-GCGCGCTCGAGTTGATTTAACGGCTGCTGTAATG-3' (SEQ ID NO:11). The amplified fragment was digested and cloned into the *Nco*I and *Xho*I sites of pMEB16, which contains the sequence encoding the N-terminal 123 amino acid residues of the *Ssp* DnaE intein. The resulting vector pEA(N) expresses a fusion protein composed of the ALSII_m N-terminal fragment and the DnaE N-terminal

fragment (ALSII_m(N)-IN_n). The ALS II C-terminal fragment was amplified using primers 5'-GCGCGACCGGTTGTGACTGGCA GCAACACTGC-3' (SEQ ID NO:12) and 5'-GGGGGGCTGCAGTCA TGATAATTTCTCCAAC-3' (SEQ ID NO:13). The fragment was
5 digested with *Age*I and *Pst*I and then cloned into the *Age*I and *Pst*I sites of pMEB9. The resulting plasmid pEA(C) expresses a fusion protein composed of the *Ssp* DnaE intein C-terminal fragment and the ALSII C-terminal fragment (ALSII_m(C)-IN_C). A 1kb *Xba*I-*Pst*I fragment containing
10 ALSII_m(C)-IN_C fusion gene was subcloned from pEA(C) into the *Xba*I and *Pst*I sites of pKEB1 plasmid to produce a kanamycin resistant expression vector pKEC3.

When pEA(N) and pKEC3 were co-expressed in *E. coli*
15 ER2744, it was predicted that *trans*-splicing of the two fusion proteins would result in ligation of the two split halves of the *E. coli* ALSII_m, with 14 amino acids present at the ligation junction.

20 6. Characterization of protein *trans*-splicing activity

To determine whether ALSII-DnaE intein fusion proteins are able to *trans*-splice in *E. coli* cells to produce ALSII_m-14,
25 western blots were performed using rabbit antiserum specifically against either the N- or C-terminal fragment of ALSII.

Two rabbit antisera were raised against peptides derived from the N-terminal and C-terminal regions of ALSII, respectively (COVANCE). These two peptides are 1) NH₂-CAQ WVVHALRAQGVNTVFGYG-COOH (SEQ ID NO:8) derived from the ALSII N-terminal sequence (amino acid residues Ala4 to Tyr23) and 2) NH₂-CVWPLVPPGASNSEMLEKLS-COOH (SEQ ID NO:9) derived from the ALSII C-terminal sequence (amino acid residues Val 530V to Ser548). A single bacterial colony was inoculated in LB medium supplemented with 100 µg/ml of ampicillin for 4 hrs at 37°C. Then it was induced by addition of IPTG to 0.3 mM final concentration. Cells were further cultured for 2-16 hours at 15°C. 20 µl of cell culture was removed, mixed with 3XSDS loading buffer (New England Biolabs, Beverly, MA), boiled for 5 minutes and 2 µl was loaded to 12% Tris-glycine gel (Novex, San Diego, CA). Subsequently proteins were transferred to a nitrocellulose membrane and blocked with 5% dry milk for one hour at room temperature (Sambrook, *et al.*, Molecular Cloning, (1989)). Immunoblotting was performed using antiserum (1:20000 dilution) overnight at 4°C in the presence of 1% dry milk. Blots were then washed three times for 15 minutes each and incubated with 1:10000 diluted HRP-conjugated anti-rabbit secondary antibody for 1 hour at room temperature. The reactions were visualized with Chemiluminescent Western Detection kit (New England Biolabs, Beverly, MA).

In control cells cultured at 15°C, expression of ALSII (Figure 8A, 8B & 8C, lane 2) was recognized specifically by

both antibodies. In cells bearing a single ALSII-intein fusion vector and another control vector to confer both ampicillin and kanamycin resistance only ALS(N)-IN_n or ALS(C)-IN_c protein was detected by anti-ALS(N) or anti-ALS(C) serum (Figure 8A, lane 3, Figure 8B, lane 4). When ALS(N)-IN_n and ALS(C)-IN_c were co-expressed a 60 kD band, as expected for the spliced product ALSII_m-14, reacted with antibodies raised against the N-terminus and C-terminus of ALSII (Figure 8A & 8B, lane 5). This band of ALSII_m-14, as predicted, exhibited a slightly higher molecular weight than native ALSII. The data indicated that trans-splicing occurred between the two ALSII-intein fusion proteins. A non-specific protein reacting with anti-ALS(N) was observed (Figure 8A and Figure 8C, lane 1 to lane 5).

Trans-splicing activity of the *Ssp* DnaE intein was previously shown to be temperature sensitive (Evans *et al.*, *J. Biol. Chem.* 275:9091-9094 (2000)). The temperature sensitivity of *trans*-splicing of the ALSII-*Ssp* DnaE intein proteins were examined by western blot analysis using an antiserum against ALSII N-terminal fragment (Figure 8C). Cells were transformed by plasmids expressing ALSII, or both ALSII_m(N)-IN_n and ALSII_m(C)-IN_c. Expression of the ALSII proteins were induced at 37°C for 3 hours. Co-expression of ALSII_m(N)-IN_n and ALSII_m(C)-IN_c was induced at 37°C for 3 hours, 30°C for 3 hours, 25°C for 6 hours, or 15°C for 16 hours. Cell extracts were treated with SDS sample buffer and denatured at 95°C to 100°C for 5 minutes and then

subjected to electrophoresis on a 12% SDS-PAGE. A western blot was probed using an antiserum raised against the ALSII N-terminal fragment. Figure 8C includes the following samples: cells with no ALSII (lane 1, control), ALSII (lane 2), ALSII_m(N)-IN_n and ALSII_m(C)-IN_c (lane 3 to lane 6). The cell culture temperature are 37°C for lane 1 to lane 3, 30°C for lane 4, 25°C for lane 5, and 15°C for lane 6.

In cells grown at 37°C, ALSII_m-14 was not detectable (Figure 8C, lane 3). However, in cells cultured at 30°C, the spliced product was observed with a significant amount of N-terminal fusion protein accumulation (Figure 8C, lane 4). In cells cultured at 25°C and 15°C (Figure 8C, lane 5 and 6), only the spliced product was detected, indicating a complete conversion of the N-terminal fusion protein to the spliced product. The ALSII_m(C)-IN_c protein was produced in excess under all the expression conditions. The data demonstrated that the *Ssp* DnaE intein was capable of mediating *trans*-splicing of the N- and C-terminal ALSII_m protein segments to form ALSII_m-14. The splicing reaction was inhibited when the experiment was conducted at 37°C. Splicing appeared to be more efficient when cells were cultured at 15°C -25°C rather than at 30°C.

7. Herbicide resistance in cells bearing the split ALS gene

The next step was to determine whether the spliced product, as the result of *trans*-splicing of the ALSII_m

(ALSII_m(N)-IN_n and ALSII_m(C)-IN_c) fusion proteins, would render *E. coli* ER2744 resistant to valine and SM. The first experiment was to test the effect of co-expression of ALSII_m(N)-IN_n and ALSII_m(C)-IN_c fusion proteins on cell growth in valine saturated M9 minimum medium. In a plating assay (see Section 4.), all transformed cells grew well on M9 medium in the absence of valine (Figure 9A-a). However, only ALSII and its herbicide resistant mutant ALSII_m rescued the cells growth at both 30°C and 37°C (Figure 9A-b, 9A-c) in the presence of valine. Significantly, co-expression of ALSII(N)-IN_n and ALSII(C)-IN_c rescued cell growth at 30°C (Figure 9A-c) or lower temperatures (data not shown) from a valine plate. Furthermore, expression of ALSII_m or ALSII_m(N)-IN_n and ALSII_m(C)-IN_c rescued cells from additional herbicide inhibition (Figure 9A-d). Moreover, transformation of wild type ALSII could not rescue cell growth from herbicide inhibition (Figure 9A-d). The control cells which expressed either ALSII(N)-IN_n or ALSII(C)-IN_c alone did not grow on a valine plate (Figure 9A-b, 9A-c); neither did the co-expression of native ALSII N- and C-terminal segments which were not fused to the intein (Figure 9A-b, 9A-c). The data indicates that co-expression of ALSII_m(N)-IN_n and ALSII_m(C)-IN_c fragments is required for *trans*-splicing and generating a functional ALSII, which can rescue cell growth from valine and herbicide inhibition.

A quantitative liquid culture assay was performed to verify the results obtained from the plating assay. The liquid assay was performed as follows. A single colony was used to

inoculate LB medium supplemented with kanamycin and ampicillin at 37°C for 4 hours. Expression was induced by 0.3 mM IPTG and cell cultures were shifted to 30°C for another 2hrs. Then, 200 µL of equivalent OD₆₀₀ 8.0 was spun down, washed one time with M9 medium and resuspended in 200 µL M9 medium. 40 µL was aliquoted into 2ml of appropriate culture medium and grown for 24-72 hours before its OD₆₀₀ was measured. The concentration for valine is 100 µg/ml and for SM is 50 µg/ml. At 30°C, all transformed cells grew equally well in M9 minimum medium (Figure 9B). In valine saturated M9 medium, wild type ALS allowed cells to grow, but no growth was observed when SM was added. However, the expression of ALSII_m or co-expression of ALSII_m(N)-IN_n and ALSII_m(C)-IN_c allowed cells to grow in valine M9 medium, as well as medium containing SM. In control experiments, ALSII_m(N)-IN_n or ALSII_m(C)-IN_c alone or co-expression of ALSII_m N- and C-terminus not fused to the intein, did not rescue cell growth in valine containing medium. This data is in agreement with that from the plating assay. To further compare the growth kinetics for *trans*-splicing mediated cell growth to wild type ALSII mediated cell growth, a time course study was performed (Figure 9C). Data showed that ALSII expressing cells have the fastest growth rate followed by ALSII_m expressing cells. ALS(N)-IN_n and ALS(C)-IN_c transformed cells have slower growth rates compared to ALSII wild type expressing cells, but not significantly less than the growth rate of ALSII_m expressing cells. Cells expressing split ALSII, with no fusion to the intein, have very slow

growth. Therefore, from the plating and liquid assay we have demonstrated that the *Ssp* DnaE can mediate ALSII *trans*-splicing, which results in a functionally herbicide-resistant ALSII_m-14 *in vivo*.

5

In conclusion, the data indicated that the two ALS-intein fusion proteins, produced from two different loci, underwent *trans*-splicing in a temperature-dependent manner to form a full length, functional ALSII_m protein. *E. coli* host cells possessing both ALSII_m fusion gene fragments showed the herbicide resistance phenotype.

10

EXAMPLE II

15

Trans-splicing of a maize Acetolactate Synthase in *E. coli*

In this Example, we demonstrate a method to produce a full length maize acetolactate synthase by protein *trans*-splicing in *E. coli*. We demonstrate how to choose a split site in the maize ALS gene based on the sequence homology of the maize ALS gene and its *E. coli* counterpart, the ALSII gene. We show that, when the split maize ALS-intein fusion genes were co-expressed, the two fusion proteins underwent *trans*-splicing to produce a protein product of expected size for the mature maize ALS protein.

20

25

1. Selection of split site.

It is important to demonstrate the *trans*-splicing of other herbicide resistant genes, such as the maize acetolactate synthase (cALS) gene, of which the herbicide resistant mutant form has been utilized to genetically modify plants (Bernasconi *et al.*, *J. Biol. Chem.* 270:17381-17385 (1995)). One preferred method for the identification of a suitable split site within any gene is to analyze the *trans*-splicing activity of a homologous gene from a different organism. We have described in Example I, that *E. coli* ALSII gene, after being split between Q327 and C328, can be reconstituted by the *trans*-splicing activity of the *Ssp* DnaE intein *in vivo*. Sequence alignment between *E. coli* ALSII and maize ALS was conducted to search for the region in the maize ALS gene corresponding to the split site of the *E. coli* ALSII gene. The result suggests that Serine397 and Threonine398 align with the split site (Glutamine327 and Cysteine328) of *E. coli* ALSII. Splitting the maize ALS between Serine397 and Threonine398, as indicated by a star (Figure 6), may yield two maize ALS-intein fusion proteins which would be capable of proficient splicing.

2. Cloning of the maize ALS gene

Reverse transcriptase polymerase chain reaction (RT-PCR) was carried out to clone the maize ALS cDNA.

Total RNA was isolated from corn leaves using the RNAqueous kit (Ambion, Inc., Texas). The RNA was then used

for first strand cDNA synthesis using the reverse primer 3-3 (5'-AT CAGTACACAGTCCTGCCATC-3' (SEQ ID NO:14)) and Superscript Reverse Transcriptase (LTI-GIBCOBRL, Rockville, MD). The first strand cDNAs were then treated with RNaseH (LTI-GIBCO BRL, Rockville, MD) before being used as a
5 template in a PCR reaction. The PCR reaction was carried out using Expand Long Template PCR system (Boehringer Mannheim, Germany). The primers used in this reaction were Reverse Primer 3-3 and cALS 5-4 primer (5' GAGACAGCCGCCG
10 CAACCAT-3' (SEQ ID NO:15)).

An aliquot of the PCR product was electrophoresed on an agarose gel and a band of approximately 2 kb was observed. This fragment was cloned into the TOPO 2.1 vector (Invitrogen, San Diego, CA, manufacturer's protocol) to make
15 pCALS1. The sequence of pCALS1 was confirmed using M13 forward and reverse primers.

3. Construction of the maize ALS-intein fusions

20

The DNA encoding for the N-terminal 397 amino acid residues of the maize ALS gene was amplified by PCR using forward primer 5'-GGGCCCATATGGCCACCGCCGCCGCCGCG-3' (SEQ ID NO:16), reverse primer 5'-GGGCCCTCGAGGCTTCCTTC
25 AAGAAGAGC-3' (SEQ ID NO:17), and the template pCALS1 (Sambrook *et al.*, *Molecular Cloning*, (1989)). A 1.2 kb PCR product was cloned into TOPO-blunt vector (Invitrogen, San Diego, CA manufacturer's protocol), resulting in TOPO-cALS(N).

Then TOPO-cALS(N) was digested with *Nde*I and *Xho*I. A 1.2 kb digested DNA fragment was recovered from low melting agarose gel and fused in-frame to the DNA sequence encoding the N-terminal 123 amino acids of the *Ssp* DnaE intein, resulting in a vector (MEB10-cALS(N) which expresses the N-terminal cALS-intein fusion protein, cALS(N)-IN-n. A DNA fragment encoding for the C-terminal 241 amino acid residues of the maize ALS gene was PCR amplified using forward primer 5'-GGGCCACCGGTACATCAAAGAAGAGCTTG-3' (SEQ ID NO:18), reverse primer 5'-GGGGCTGCATTTCAGTACACAGTCCTGCATC-3' (SEQ ID NO:19), and the template pCALS4. A 0.8 kb PCR product was cloned into a TOPO-blunt vector (see protocol above), TOPO-cALS(N). TOPO-cALS(N) was then digested with *Age*I and *Pst*I. A 700 bp DNA fragment was recovered from low melting agarose gel and was fused in frame to the DNA encoding the C-terminal 36 amino acids of the *Ssp* DnaE intein, resulting in a vector MEB9-cALS(C). MEB9-cALS(C) was further cut by *Xba*I and *Pst*I and released a 1 kb fragment. This 1 kb fragment was cloned into pKEB1 vector to create a kanamycin resistant expression vector for the cALS-intein C-terminal fusion protein, cALS(C)-IN_C. The same extra 7 amino acids, NH₂-LEKFAEY-COOH (SEQ ID NO:20) and NH₂-CFNKSTG-COOH (SEQ ID NO:21) were also present at the junctions of the N- and C-terminal cALS-intein fusion proteins, respectively.

4. Trans-splicing of the maize ALS-intein fusion proteins

Both ALS-intein fusion fragments, cALS(N)-IN_n and cALS(C)-IN_c, described in Section 3, were co-expressed in *E. coli* ER2744 under the same conditions as described in Example I, Section 6. A western blot was performed to detect the *trans*-splicing product (Method, see Example I, Section 6). On the blot, a fragment of 69 kD, which corresponds to the size of the wild type cALS (Figure 10A, and Figure 10B, lane 2), was detected in both fusion proteins expressed cells and was recognized by rabbit antisera specifically raised against two peptides derived from N- and C-terminal sequence of maize ALS (Figure 10A and Figure 10B, lane 5). A non-specific protein reacting with antiserum against N-terminal of maize ALS was observed (Figure 10A, lane 1 to lane 5). The peptides used to raise antibodies are 1) ALS-N peptide corresponding to the sequence from Lys66 to Ala85, NH₂-CKGADILVESLERCGVRDVFA-COOH (SEQ ID NO:22), and 2) ALS-C peptide corresponding to the sequence from Ile619 to Tyr638, NH₂-CI PSGGAFKDMILDGDGRTVY-COOH (SEQ ID NO:23). The full length cALS species was not detected in cells expressing either N- or C-terminal fusion protein (Figure 10A and Figure 10B, lane 3 and lane 4). This demonstrated that split maize ALS, like *E. coli* ALSII, when fused with the *Ssp* DnaE intein, was also able to perform *trans*-splicing to produce the full length ALS.

In conclusion, the maize ALS gene was split by the *Ssp* DnaE intein and cloned into two separate plasmid vectors. When both the fusion gene vectors were introduced into the same host cell and co-expressed, the two fusion proteins
5 underwent *trans*-splicing to produce a full length cALS. Although a functional assay is needed to determine the activity of the spliced maize ALS protein in plants, it does raise the possibility of successfully splitting a plant herbicide
10 resistant or disease resistant genes into two inactive gene segments. These two gene fragments can be confined into two separate cellular compartments, such as the chloroplast and nucleus, or two separate loci on the chromosomes, or two separate DNA vector. This novel mode of gene expression
15 may greatly lessen the chance of spreading an intact active transgene into other species.

EXAMPLE III

The present Example details the feasibility of splitting
20 the *aroA* gene and regenerating the desired protein activity using an intein. The experiment consisted of dividing the gene encoding the mutant *aroA* gene at various positions and fusing the gene encoding the N-terminal splicing domain of
the *Ssp* DnaE intein (IN_n) to the gene encoding the N-terminal
25 fragment of the EPSPS protein. Concurrently, the gene encoding the C-terminal splicing domain of the *Ssp* DnaE intein (IN_c) was fused to the gene encoding the C-terminal
fragment of the EPSPS protein. When the fusion genes were

placed on to two separate plasmids and co-transformed and co-expressed in the same bacterial cell it was demonstrated that those bacterial cells were resistant to the herbicide glyphosate.

The cloning of *Salmonella typhimurium aroA* gene that confers resistance to glyphosate

1. Creation of plasmid pEPS#1

The *Salmonella typhimurium aroA* gene with the C301 to T mutation was acquired from the American Type Culture Center in the form of a cosmid in the bacteria *Salmonella choleraesuis* subsp *choleraesuis* (ATCC No. 39256). The modified *aroA* gene was amplified from the cosmid by the polymerase chain reaction using primers EPSP#1 (5'-GGATCCTAAGAAGGAGATATACCCATGGAATCCCTGACGTTACA-3' (SEQ ID NO:24)) and EPSP#2 (5'-GTCGACGCTCTCCTGCAGTTAGGCAGGC GTACTCATTC-3' (SEQ ID NO:25)). The PCR product was inserted into the *StuI* site of the plasmid LITMUS 28 (New England Biolabs, Inc., Beverly, MA). Following transformation and plasmid preparation, sequencing revealed an unexpected mutation (C103 to G) which was reverted using Stratagene's (La Jolla, CA) Quick Change Site Directed Mutagenesis Kit and primers EPSP#10 (5'-GCTTTGCTCCTGGCGGCTTTACCTTGTGGT AAAACCGC-3' (SEQ ID NO:26)) and EPSP#11 (5'-GCGGTTTTAC CACAAGGTAAAGCCGCCAGGAGCAAAGC-3' (SEQ ID NO:27)). Sequencing of DNA from the resulting colonies revealed that the unexpected mutation had been reverted to the expected

C. This plasmid was termed pEPS#8 and used as the acceptor plasmid in the subsequent transposition linker scanning reactions.

5 **2. Description of ER2799, an *E. coli* strain used to test the *aroA* gene constructs**

 An *E. coli* strain that has the *aroA* gene deleted from its chromosome was acquired from the Yale *E. coli* stock center
10 (*E. coli* strain AB2829, CGSC#2829, ID#8215). This strain was made hsdR- and named ER2799. Because ER2799 lacks the *aroA* gene, which is necessary for aromatic amino acid synthesis, it does not grow on M9 minimal media. This strain is used to test the various *aroA* gene constructs to see if the
15 new *aroA* gene can rescue the bacteria and allow growth on minimal media either in the presence or absence of glyphosate.

20 **3. Finding a site to split the *aroA* target gene by transposon based linker scanning**

 The first step in performing this experiment was to determine the sites in the 5-enolpyruvyl-3-phosphoshikimate synthetase (EPSPS) protein which could allow insertion of an
25 intein *in cis*. *In cis* refers to the fact that the complete intein is inserted into the complete EPSPS protein. However, it was not known which portions of the EPSPS protein itself would be tolerant to extra amino acid residues. So to determine where the EPSPS protein could tolerate amino acid insertions a new
30 technology, the GPS®-LS kit (available from New England

Biolabs, Inc., Beverly, MA), was used to randomly insert 5 amino acid residues throughout the EPSPS protein sequence. An expression plasmid library was constructed with the EPSPS gene with the randomly inserted 5 amino acids. This library was transformed into *E. coli* strain ER2799 and applied to plates containing M9 minimal media. ER2799 lacks the *aroA* gene and will not grow on M9 minimal plates unless an active EPSPS gene is supplied by plasmid transformation. The ER2799 *E. coli* that grew following transformation with the library should contain an EPSPS protein that is active with the 5 amino acid insertion. These were sequenced to determine the position of the 5 amino acid insertion and 42 unique sites were discovered in the EPSPS protein that allowed growth of ER2799 on M9 minimal plates (Figure 15). Furthermore, another 19 unique sites were found that did not tolerate a 5 amino acid insertion (Figure 16).

4. Transposition Reaction

The reaction was performed by adding 6 μ l of 20 ng/ μ l pEPS#8 (target DNA), 1.5 μ l of 20 ng/ μ l *Pme*I donor DNA, 3 μ l of distilled water, 3 μ l of 10X GPS®-LS buffer and 1.5 μ l of Tn*ABC and mixing for 15 min at 37°C. 1 μ l of Start Solution was added and the reaction incubated at 37°C for 1 hour and 20 min. The reaction was stopped by heat inactivation for 15 min at 75°C. Following cooling the reaction mixture to room temperature and dialysis against water for 2 hours the reaction mixture was transformed into freshly-made ER2685.

(fhuA2 glnV44 el4- rfbD1? relA1? endA1 spoT1? thi-1 Δ (mcrC-mrr)114::IS10 Δ (lacI-lacA)200 F'proA+B+lacIq D1 (lacZ)M15 zsf:Tn10 (TetR)) cells by electroporation. The cells were incubated for 1 hour at 37°C and then plated onto LB plates containing ampicillin and kanamycin. Cell growth was allowed to proceed at 37°C overnight. It was discovered that 10 μ l of reaction mixture gave over 10,000 colonies (enough to cover all possible transposon insertion sites, 2840 sites in pEPS#8, 3.3 times) following transformation.

5. Isolating the DNA fragment (3.0kb) containing the EPSPS gene plus transposon

All the transformants from the transposition reaction were recovered using LB medium and 66% of the cells were saved at -70°C by adding 20% glycerol. The rest were grown in 500 ml of LB liquid medium containing 100 μ g/ml ampicillin, and 50 μ g/ml kanamycin at 37°C overnight. The cells were harvested by centrifugation and the plasmid DNA was purified (508 μ g total) using a Qiagen Midi kit (Qiagen, Studio City, CA). The 3.0kb *aroA* gene-Transposon DNA fragment was released by digesting the DNA(58 μ g) with *Pst*I, *Nco*I and *Ahd*I and isolated by gel-purification using agarase following ethanol precipitation (4 μ g DNA was recovered).

6. Cloning the *aroA* gene-Transposon 3.0 kb fragment into the pCYB3 vector

The gel-purified 3.0 kb *aroA* gene-Transposon DNA fragment was ligated into the *Nco*I to *Pst*I sites of pCYB3

(5.2kb), and transformed into ER2685 by electroporation after drop dialysis for 2 hours. The electroporated cells were incubated for 1 hour in LB medium. 250 μ l of this cell suspension was plated onto LB plates containing 100 μ g/mL ampicillin and 50 μ g/mL kanamycin while another 5.5 ml was inoculated into 1 liter of LB liquid medium with 100 μ g/mL ampicillin and 50 μ g/mL kanamycin and grown at 37°C overnight. The plasmid DNA library containing the transposon within the *aroA* gene was isolated by Qiagen (Studio City, CA) Midi kit (750 μ g).

7. Screening the library EPSPS protein that is active with the 5 amino acid linker

105 μ g of the library DNA was digested with *PmeI* to remove the transposon from the *aroA* gene. This leaves 15 bases (or 5 amino acid residues) at the transposon insertion site. A 7 kb fragment was recovered (in a final volume of 400 μ l EB), self-ligated (86 μ l out of 400 μ l 7 kb fragment in a 100 μ l rxn), transformed (30 μ l of the 100 rxn) into *E. coli* strain ER2799 and plated onto both LB and M9 minimal plates, each containing 100 μ g/mL ampicillin in the presence of 0.3 mM IPTG. Following incubation at 37°C overnight *ca.* 20% of the original cells survived on M9 minimal plates as compared to the LB plates. Individual colonies that grew on M9 minimal media plates were analyzed by *DraI* digestion and DNA sequencing to confirm the site of linker insertion site into the *aroA* gene.

42 different insertion sites were identified among 72 active individual clones that can tolerate 5 amino acid residues inserted into the *aroA* gene and 19 different insertion sites were identified among 39 inactive clones that can not grow on M9 minimal media selection plates (see Figure 15 and Figure 16). Plasmids pCE-5-22, pCE-5-21, pCE-5-35 and pCE-5-23 were the active clones that have 5 amino acid residues incorporated into the EPSPS protein (*aroA* gene product) at positions 182, 215, 235 and 267, respectively. These four sites were chosen for further studies.

Construction of *Ssp* DnaE Cis- and Trans-splicing vectors

1. Creation of vectors pCE182DnaE, pCE215DnaE, pCE235DnaE, and pCE267DnaE for Cis-Splicing

This involved inserting an intein into the sites in the target protein that were discovered to tolerate 5 amino acid insertions.

Four sites were chosen for further study (positions 182, 215, 235, and 267). The full length *Ssp* DnaE intein was inserted into these sites and the EPSPS-intein fusion was tested for its ability to permit ER2799 cells to grow on M9 minimal plates. All four sites were found to grow on M9 plates, indicating that the EPSPS protein could tolerate the intein inserted at these positions (see Figure 11 and Figure 14).

CE182 or CE215, which was the linear DNA of pCE-5-22 or pCE-5-21 with the exception that the five amino acid linker at 182 or 215 has been removed, was generated by polymerase chain reaction (PCR) from templates pCE-5-22 or pCE-5-21 using primers 5'-GCCCCTAAAGACACAATTATTCGCG-3' (SEQ ID NO:28) and 5'-CAGCGGCGCCGTCATCAGCAGAGCG-3' (SEQ ID NO:29) for CE182 or 5'-GCGAACCACCACTACCAACAATTG-3' (SEQ ID NO:30) and 5'-TATCTCCACGCCAAAGGTTTTTCATT-3' (SEQ ID NO:31) for CE215. The *Ssp* DnaE intein gene containing two native N-extein residues and three native C-extein residues was amplified by PCR from pMEB8 (Evans, et al., *J. Biol. Chem.*, 275:9091 (2000)) using primers 5'-GAATATGCTGTCTTTTGGT-3' (SEQ ID NO:32) and 5'-GTTAAAGCAGTTAGCAGCGAT-3' (SEQ ID NO:33). The resultant PCR fragment was phosphorylated with T4 polynucleotide kinase, purified by QIAquick column (Qiagen, Inc., Studio City, CA) and ligated into CE182 or CE215 to generate pCE182DnaE or pCE215DnaE, respectively.

The *Ssp* DnaE intein gene containing four native N-extein residues and three native C-extein residues was amplified by PCR from pMEB8 using primers 5'-TGCTGAATATTGCTGTCTTTTGG-3' (SEQ ID NO:34) and 5'-CCGTTAAAGCAGTTAGCAGCGATAGC-3' (SEQ ID NO:35). The resultant PCR fragment was purified by QIAquick column (Qiagen Inc., Studio City, CA) and ligated into the gel-purified, *Pme*I cut pCE-5-35 or pCE-5-23 vector DNA to generate pCE235DnaE or pCE267DnaE, respectively.

2. Creation of vectors p215EN2/pEPS#28 and p235EN2/pEPS#29 for Trans-Splicing:

Two plasmids were constructed with compatible origins of replication. The N-terminus of the appropriate EPSPS protein was fused to the N-terminus of the N-terminal *Ssp* DnaE splicing domain (IN_n) and inserted into one plasmid. The remaining C-terminal portion of the appropriate EPSPS protein was fused to the C-terminus of the C-terminal splicing domain of the *Ssp* DnaE intein (IN_c). This fusion was inserted into the second plasmid. The plasmids were co-transfected into ER2799 by electroporation. Expression of the fusion protein was under the control of an IPTG inducible pTac promoter. The transformed cells grew on M9 minimal plates, liquid M9 minimal media, or liquid M9 minimal media supplemented with glyphosate (Figures 11, 12, 13 and 14). This indicated that the protein halves could generate an active EPSPS protein when co-expressed in the same cell.

The 0.6 kilobase *Xho*I to *Pst*I fragment of pMEB4 was gel-purified using the QIAquick extraction kit and ligated into the *Xho*I to *Pst*I sites in the pCYB3 (New England Biolabs, Inc., Beverly, MA) vector to generate pCEN1. The *Nco*I site between the *Ssp* DnaE intein and the chitin-binding domain (CBD) was removed by *Pac*I and *Sap*I digestion of pCEN2 followed by T4 DNA polymerase treatment and self-ligation to generate plasmid pCEN2. This vector contains the N-terminal 123 amino acid residues of the *Ssp* DnaE intein (IN_n) under

the control of pTac promoter and confers resistance to ampicillin.

5 p215EN2 or p235EN2 were constructed by ligating the *Nco*I to *Kpn*I fragment of pCE215DnaE or pCE235DnaE into the same sites of pCEN2. p215EN2 or p235EN2 has the N-terminus of EPSPS (residues 1-215 for p215EN2, 1-235 for p235) fused to the IN_n.

10 The *Nco*I to *Fsp*I fragment of pCYB3 was ligated into the *Nco*I to *Dra*I sites of pKEB1 to generate pKEB12 (NEB#1282). A sample of pKEB12 plasmid transformed in *E. coli* strain ER2566 has been deposited under the terms and conditions of the Budapest Treaty with the American Type Culture
15 Collection on May ____, 2000 and received ATCC Patent Accession No. _____. This vector has the C-terminal 36 amino acid residues of the *Ssp* DnaE intein (IN_n) fused to CBD and confers resistance to kanamycin.

20 pEPS#28 and pEPS#29 were constructed by ligating the *Bgl*II to *Pst*I fragment of pCE215DnaE and pCE235DnaE into the same sites of pKEB12. pEPS#28 or pEPS#29 has the C-terminus of EPSPS (residues 216-427 for pEPS#28, 236-427 for pEPS#29) replacing the CBD in pKEB12 and attached to
25 the C-terminus of IN_C.

3. Creation of the EPSPS complementary construct pEPS#34 and pEPS#36.

When the EPSPS protein fragments, lacking the intein domains, were co-expressed in ER2799 cells, the cells failed to grow on M9 minimal plates, liquid M9 minimal media, or liquid M9 minimal media supplemented with glyphosate (Figure 12 and Figure 13). This indicated that EPSPS activity was absolutely dependent on the presence of both intein halves.

DNA encoding the N-terminus of the EPSPS protein, residues 1-235, (EPS235N) was amplified by PCR from pCE235DnaE using primers 5'-GGATCCTAAGAAGGAGATATACCC ATGGAATCCCTGACGTTACA-3' (SEQ ID NO:36) and 5'-GATATC CTGCAGTTAACCTGGAGAGTGATACTGTTGACC-3' (SEQ ID NO:37). The resultant PCR product was purified using a QIAquick PCR kit, digested with *Nco*I and *Pst*I, purified from an agarose gel using the QIAquick extraction kit and ligated into the *Nco*I to *Pst*I sites of plasmid pCYB3 to generate pEPS#34.

Plasmid pEPS#36 was created by amplifying DNA encoding the C-terminus of EPSPS, residues 236-427, (EPS235C) by PCR from pC+E2 using primers 5'-GATATCCCATG GGACGCTATCTGGTCGAGGGCGATG-3' (SEQ ID NO:38) and 5'-GT CGACGCTCTCCTGCAGTTAGGCAGGCGTACTCATTC-3' (SEQ ID NO:39). The resultant PCR product was purified using the QIAquick PCR kit, digested with *Nco*I and *Pst*I, purified from

agarose gel and ligated into the *Nco*I to *Pst*I sites of plasmid pKEB12. Two extra residues Met-Gly were also incorporated at the N-terminus of EPS235C due to the *Nco*I site for cloning.

5 **4. Creation of Vectors Containing the Cis or Trans
 "dead" *Ssp* DnaE intein at position 235
 (pEPS#31, pEPS#33, pEPS#37).**

10 Interestingly, trans-splicing was not required for activity,
because if three of the most highly conserved catalytic
residues of the *Ssp* DnaE intein were changed to alanine the
co-transformed ER2799 cells still grew. This event
demonstrates that the intein can act as an affinity domain to
bring the two EPSPS intein fragments together (Figure 12 and
15 Figure 13).

20 The *Ssp* DnaE intein gene containing four native N-
extein residues and three native C-extein residues was
amplified by PCR from pMEB8 using primers 5'-TGCTGAATATGC
GCTGTCTTTTGGTACCGAA-3' (SEQ ID NO:40) and 5'-CCGTTAAA
CGCCGCAGCAGCGATAGCGCC-3' (SEQ ID NO:41). The resultant
PCR fragment was purified by QIAquick column (Qiagen Inc.,
Studio City, CA) and ligated into the *Pme*I site of plasmid pCE-
5-35 to generate pEPS#31. This *Ssp* DnaE intein contains
25 three mutations, Cys1 → Ala/Cys+1 → Ala/Asn159 → Ala, in
the catalytic residues that eliminates its splicing activity.

5. Methods of Assaying EPSPS Activity

Plating assay for EPSPS activity. The presence of a functional EPSPS protein could be determined *in vivo* using *E. coli* strain ER2799, which lacks an endogenously active EPSPS (see above). ER2799 cells alone fail to grow on M9 minimal plates (supplemented with 0.3 mM IPTG). In the following description when M9 minimal plates are mentioned they also contain 0.3 mM IPTG. Plasmid pC+E2, which contains the full length wild type EPSPS gene with a C301 to T mutation, is able to rescue growth of ER2799 on the M9 minimal plates when introduced by transformation.

Assaying the *Ssp* DnaE cis-splicing constructs. Plasmids pCE182DnaE, pCE215DnaE, pCE235DnaE, pCE267DnaE (0.05 µg of each) were transformed into *E. coli* ER2799 cells by electroporation (Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory, NY: Cold Spring Harbor Laboratory Press (1989)), see Fig. 11. 0.8 mL of LB media was added to the transformed cells and these were incubated at 37°C for 1 hour with shaking. 200 µL of this solution was plated onto either LB or M9 minimal plates supplemented with 0.1 mg/mL ampicillin (Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory, NY: Cold Spring Harbor Laboratory Press (1989)). The plates were incubated for varying length of time and at various temperatures. The most commonly used being overnight at 37°C.

Assaying the *Ssp* DnaE trans-splicing constructs. The activity of each EPSPS trans construct was assayed by co-transforming the constructs to be tested into ER2799 and plating on either an M9 minimal plate, containing 0.3 mM IPTG, or an LB plate in which both were supplemented with 0.1 mg/mL ampicillin and 0.05 mg/mL kanamycin. In cases where only one plasmid contained the EPSPS gene or a portion of the EPSPS gene the complementary antibiotic resistance was supplied by co-transforming the *E. coli* with either pCYB3 or pKYB1 (New England Biolabs, Beverly, MA), which has no EPSPS gene present.

The plasmids used were: pC+E2, p215EN2, p235EN2, pEPS#28, pEPS#29, pEPS#33, pEPS#37, pEPS#34, and pEPS#36. These plasmids were co-transformed (Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory, NY: Cold Spring Harbor Laboratory Press (1989)) using 0.1 µg of the appropriate plasmids, in various combinations, into ER2799 *E. coli* cells, and plated on both LB plates and M9 minimal media plates, each containing 100 µg/mL ampicillin and 50 µg/mL kanamycin. The M9 minimal plate also contained 0.3 mM IPTG. Individual clones were picked from each LB plate and stripped on one M9 minimal media selection plate following incubation at 37°C overnight or RT for 2-3 days. The combinations used were: WT, pC+E2 and pKYB1 (New England Biolabs, Beverly, MA); 215NC, p215EN2 and pEPS#28; 215C, pEPS#28 and pCYB3;

235NC-Dead, pEPS#33 and pEPS#37; 235NC, p235EN2 and pEPS#29; 235N, p235EN2 and pKYB1; 235C, pEPS#29 and pCYB3; 235N-215C, p235EN2 and pEPS#28; and 235 complement, pEPS#34 and pEPS#36 (see Figure 12).

5

Determination of ER2799 growth in liquid culture in the presence or absence of glyphosate. The testing of glyphosate resistance for the 235 trans constructs was made using plasmid combinations as follows; WT, pC+E2 and pKYB1; 235NC-Dead, pEPS#33 and pEPS#37; 235NC, p235EN2 and pEPS#29; 235N, p235EN2 and pKYB1; 235C, pEPS#29 and pCYB3; and 235 complement, pEPS#34 and pEPS#36. These plasmids were co-transformed into ER2799 *E. coli* cells as described above and plated onto LB plates containing 100 µg/mL ampicillin and 50 µg/mL kanamycin. As a control, pCYB3/pKYB were co-transformed into *E. coli* strain ER2744, and plated on an LB plate containing 100 µg/mL ampicillin and 50 µg/mL kanamycin. A preculture was prepared for each transformation by inoculating the fresh colony into LB medium supplemented with 100 µg/mL ampicillin and 50 µg/mL kanamycin at 30°C for overnight. Equal amounts of pre-culture (10-11µL depending on the cell density) was inoculated into freshly-made M9 minimal medium containing 100 µg/ml of ampicillin, 50 µg/ml of kanamycin and 0.3 mM IPTG in the absence or presence of different amounts of glyphosate. The growth of each construct was measured by OD at 600 nm, see Figure 13.

10

15

20

25

Growth of the cis 235 construct in M9 liquid minimal media. Two plasmid vectors one with a splicing competent *Ssp* DnaE intein (235 cis) and another with a splicing incompetent intein (235 dead), pCE235DnaE and pEPS#31, respectively, were transformed into separate ER2799 *E. coli* cells and plated on LB plates supplemented with 100 µg/mL ampicillin and 50 µg/mL kanamycin. A preculture was prepared for each transformation by inoculating the fresh colony into LB medium supplemented with 100 µg/mL ampicillin and 50 µg/mL kanamycin at 30°C for overnight. Equal amounts of pre-culture (10-11µL depending on the cell density) was inoculated into freshly-made M9 minimal medium containing 100 µg/ml of ampicillin, 50 µg/ml of kanamycin and 0.3 mM IPTG. The cell density was determined at various times using the OD at 600 nm (see Figure14).

The *Nco*I to *Kpn*I fragment of pEPS#31 was ligated into the same sites in plasmid pCEN2 to generate pEPS#33. Plasmid pEPS#37 was created by cloning the *Bgl*II to *Pst*I fragment of pEPS#31 into the same sites in plasmid pKEB12.

EXAMPLE IV

Trans-splicing of two unrelated gene products aminoglycoside-3-acetyltransferase (*aadA*) and soluble modified green fluorescent protein (*smGFP*), to give rise to a functional hybrid protein in *E.coli*.

Aminoglycoside-3-acetyltransferase gene was fused to *Ssp* DnaE intein N-fragment (IN_n). The C fragment of the *Ssp*

DnaE intein (IN_C) was fused to the smGFP gene. The fusion proteins could be translated as individual polypeptides from the respective constructs. These fusion protein coding DNA sequences were cloned into either pIH976 (Figure 17) or pAGR3 (Figure 18) plasmids. Both the plasmids (pIHaadE-N (pIH976 containing aadA and IN_n terminal) and pAGRE-CsmGFP (pAGR3 containing IN_C and smGFP)) were co-transformed in to *E. coli* (Figure 19A). The transformed *E. coli* were resistant to spectinomycin/streptomycin sulfate (Figure 19B). The cell extracts were made after 16 hrs of growth. The proteins in the extract was separated on SDS tris glycine gel and blotted on to a PVDF membrane. This membrane was probed with anti GFP monoclonal antibodies. Trans-splicing was observed in *E. coli* extracts, where both the plasmids were introduced. As a result of trans-splicing the fusion product had a molecular mass identical with the calculated cumulative mass of both the proteins (Figure 19C).

The following protocol describes the production of cassettes, pIHaadE-N (*Aminoglycoside-3-acetyltransferase* gene fused to DNA encoding IN_n), pAGRE-CsmGFP (DNA encoding IN_C was fused to *smGFP* gene), Western blotting and detection.

Polymerase chain reaction (PCR) was used for cloning of the open reading frames (ORFs) in to the desired plasmids. The reaction contains Vent[®] DNA polymerase buffer supplemented with 2 mM magnesium sulfate, 200 μM dNTPs,

1 μ M of each primer and 100 ng plasmid DNA in a total volume
of 50 μ l with 2 units of Vent[®] DNA polymerase. Between 10 to
20 rounds of amplification were carried out using a Perkin-
Elmer gene amp PCR 2400 system (Emeryville, CA). The
5 following primers used for amplification of the *aadA* gene (*aadA*
forward primer: GCCTTAATTAACCATGAGGGAAGCGGTGATCGC
CG (SEQ ID NO:47), *aadA* reverse primer: TGCGGTCTGACTTTGC
CGACTACCTTGGTGATCTC (SEQ ID NO:48). PCR products were
purified using a PCR purification kit (QIAquick PCR purification)
10 from Qiagen (Valencia, CA). Purified PCR products were
digested by *Pac* I and *Sal* I restriction enzymes and cloned in
to pNEB193 (New England Biolabs, Inc., Beverly, MA) plasmid.
The clone containing the *aadA* gene was named pNEBaad3.
Similar protocol was used for amplification and cloning of the
15 *smGFP* gene using specific primers (*smGFP* forward primer:
CCCAAGCTTGGCGCCATGAGTAAAGGAGAAGAACTTTTCAC (SEQ ID
NO:49) and *smGFP* reverse primer: GCGACCGGTTTATTTGTATAG
TTCATCCATGCCATG (SEQ ID NO:50) into pLITMUS 28 (New
England Biolabs, Inc., Beverly, MA). The clone containing the
20 *smGFP* gene was named psmGFP7. Sequences for both *aadA*
and *smGFP* genes were verified by DNA sequencing.

The Intein from the *dnaE* gene of *Synechocystis* species
PCC6803 was PCR amplified. The amino terminal part of the
25 intein (amino acids 1-123) is referred to as IN_n and the
carboxy terminal as IN_c (amino acids 124-159). Both IN_n and
IN_c fragments were cloned into pLITMUS 28 and pNEB193
respectively. The primer pairs for amplification of IN_n and IN_c

are listed (IN_n forward primer: AGGGAATTCGTCGACAAATTTG
CTGA ATATTGCCTGTCT (SEQ ID NO:51), IN_n reverse primer:
GGCCTCGAGTTATTTAATTGTCCCAGCGTCAAGTAATG (SEQ ID
NO:52), IN_c forward primer: AGCTTTGTTTAAACCATGGTTAAAG
5 TTATCGGTCGTAGATC (SEQ ID NO:53), IN_c reverse primer:
CAGCGTCGACGGCGCCGTGGGATTTGTTAAAGCAGTTAGCAGC
(SEQ ID NO:54)). The plasmids containing the IN_n and IN_c
fragments were pLitDnaE-N1 and pNEBDnaE-C2 respectively.

10 Fusion constructs of intein fragments and either *aadA* or
smGFP gene products were made in the following way:
*Bam*HI and *Sa*I fragment (800bp) from pNEBaad3 was ligated
into *Bam*HI- *Sa*I digested pLitDnaE-N1 to give rise to pAEN1.
In a similar way, the 150 bp insert (pNEBIN-c digested with
15 *Pst*I and *Ka*sI) was ligated into *Pst*I and *Ka*sI digested pLit
SmGFP5 to give rise to pGFPEC. Plasmid pAEN contains *aadA*
gene in frame with IN_n and pGFPEC contain *smGFP* gene in
frame with IN_c.

20 The fused genes were PCR amplified and cloned into
E.coli expression vectors. The inserts of pAEN and pGFPEC
were cloned into pIH976 (*Nco*I and *Sac*I site) and pAGR3
(*Eco*RI and *Sac*II sites) vectors. The primers are listed (*aadA*-
IN_n forward primers: CATGCCATGGGGGAAGCGGTGATCGC
25 CGAAG (SEQ ID NO:55), *aadA*-IN_n reverse primers: ACGCG
AGCTCTTATTTAATTGTCCCAGCGTCAAGTAATG (SEQ ID NO:56),
IN_c-*smGFP* forward primer: CGAATTCTATGGTTAAAGTTATCGG
TCGTAGATC (SEQ ID NO:57), IN_c-*smGFP* reverse primer: AG

CCCGCGGTTATTTGTATAGTTCATCCATGCCATG (SEQ ID NO:58)).

The *E. coli* expression plasmids were pIH976-aadE-N and pAGR- N_C-smGFP, under the control of Ptac promoter of the host. Either of the plasmids or both together were transformed into *E. coli* ER1992 (New England Biolabs, Inc., Beverly, MA) and plated on LB agar-Ampicillin plates as well as LB agar ampicillin and spectinomycin plates.

For Western blotting, *E. coli* cell extracts were mixed with SDS loading dye with 1 mM DTT, boiled at 95°C for 5 min and loaded on a 10-20% Tris-glycine-SDS gradient gel. The proteins were blotted on an Immobilon-P membrane and probed with an anti-GFP monoclonal antibody (Roche Molecular Biochemicals, Indianapolis, IN) followed by chemiluminescent detection of the GFP and aadA-GFP fusion protein.

EXAMPLE V

Utilization of plant promoters in *E. coli* for trans-splicing of two unrelated gene product, aminoglycoside-3-acetyltransferase (aadA) and soluble modified green fluorescent protein (smGFP) to give a functional hybrid protein.

The above DNA fragments were cloned downstream of the chloroplast specific promoter PpsbA (SEQ ID NO:59). A terminator sequence of the same gene (TpsbA (SEQ ID NO:60) was placed down stream of the cloned gene. The two genes were expressed in opposite direction to avoid read

through. The plant promoters were functional upon transformation in to *E.coli* and trans-spliced products (aadA-smGFP fusion protein, 57 kDa) were observed in Western blot assay using anti GFP antibodies. Thus chloroplast specific promoters are functional in *E.coli* and could be used for gene expression studies.

The following protocol describes the production of a *E.coli*/plant shuttle vector (pNCT114/pNCT224) that is capable of homologous recombination of a transgene(s) *in vivo*.

A shuttle vector consists of elements that will make it functional in both *E.coli* as well as plant cell. Plasmid pLITMUS28 (New England Biolabs, Inc., Beverly, MA) is the backbone for the pNCT114 and pNCT224 gene targeting vector. The vector DNA comprises, at least (1) two DNA sequence homologous to the plastid genome (also referred as targeting sequence/fragment), (2) one or more promoter element, (3) transcription terminator elements, and (4) one or more selectable/drug resistance (non-lethal) marker gene.

Promoter element (PpsbA) DNA sequences were PCR amplified from genomic DNA extracted from 7 days old tobacco seedlings using the CTAB method as described by Murray and Thompson (*Nucleic Acids Res.*, 16:4321-4325 (1980)). The primers used for amplification are listed (PpsbA forward primer: AACTGCAGGAATAGATCTACATACACCTTGG (SEQ ID NO:64), PpsbA reverse primer: CCGCTCGAGCTTAATTAAGGTAA

AATCTTGGTTTATTTAATC (SEQ ID NO:65)). Similarly the terminator sequence (TpsbA) was amplified by PCR and cloned. The primers used for amplification are listed (TpsbA forward primer: GCGACCGGTGATCCTGGCCTAGTCTATAGGAGG (SEQ ID NO:66), TpsbA reverse primer: AGGCCTAGGAGAATACT CAATCATGAATAAATGC (SEQ ID NO:67)). A vector with a psbA promoter and terminator DNA sequence allows genes to be cloned in between these for expression of the protein. The targeting DNA sequences were amplified and inserted outside of the promoter and terminator in a flanking manner (Figure 20), thus facilitating homologous recombination of the transgene at a predetermined loci. pNCT114 contains 16SrDNA-trnaV and rps7/12 targeting sequence (SEQ ID NO:61), whereas, pNCT224 contains orf228-ssb as left border and orf1244 as right border (SEQ ID NO:62). The following primers were used for PCR amplification of the targeting sequences.

Primers for pNCT114

Left border forward primer:

TTGGCGCGCTTGACGATATAGCAATTTTGCTTGG (SEQ ID NO:68)

Left border reverse primer:

TTGCGTACGATTTATCTCAGATTAGATGGTCTAG (SEQ ID NO:69)

Right border forward primer:

TTGCCTAGGCGTATTGATAATGCCGTCTTAACCAG (SEQ ID NO:70)

Right border reverse primer:

AGGGGTACCGAATTCAAGATTCTAGAGTCTAGAG (SEQ ID NO:71)

Primers for pNCT224

5 Left border forward primer:

TTGGCGCGCAATTCACCGCCGTATGGCTGACCGG (SEQ ID NO:72)

Left border reverse primer:

TTGCGTACGCCTTTGACTTAGGATTAGTCAGTTC (SEQ ID NO:73)

10

Right border forward primer:

TTGCCTAGGGTCGAGAACTCAACGCCACTATTC (SEQ ID NO:74)

Right border reverse primer:

15 AGGGGTACCATCACGATCTTATATATAAGAAGAAC (SEQ ID NO:75)

A detailed diagram for pNCT114/224 is in Figure 20A.

Both the plasmids contain two promoters and two terminator DNA fragments. For directional cloning, unique restriction enzyme sites are incorporated. Plasmid pNCT114 and pNCT224 have unique restriction enzyme sites (*PmeI*-*AgeI* and *PacI*-*XhoI* sites). Insert from plasmid pAEN (*aadA* gene in frame with IN_n) was obtained by digesting with *PacI*-*XhoI* and pGFPEC (*smGFP* in frame with IN_c) was obtained by digesting with *PmeI*-*AgeI* and ligated sequentially into pNCT114 or pNCT224. The plasmids are designated as p115ag and p225ag (Figure 21A). The plasmids were transformed into *E.coli* and selected with ampicillin and spectinomycin (Figure

20

25

21B). The cell extracts were made from overnight cultures and separated on 10-20% Tris-glycine-SDS gradient gel. The proteins were blotted on an Immobilon-P membrane and probed with an anti-GFP monoclonal antibodies (Roche Molecular Biochemicals, Indianapolis, IN) followed by chemiluminescent detection of the GFP and aadA-GFP fusion proteins (Figure 21C).

EXAMPLE VI

Cis-splicing of the EPSPS and ALS gene products in plant cytoplasm expressed from a DNA cassette integrated into molecular DNA

The introduction of DNA into plant nuclei has been achieved in many different ways, such as, electroporation, polyethylene glycol mediated, Agrobacterium mediated, microinjection and biolistic transformation. In accordance with the present invention, one should determine if the plant cytoplasm will mediate protein-splicing event in cis or trans. This will be a prerequisite for further trans-splicing technologies in plants. This technique will be useful if the target protein needs specific cytoplasmic modification for activity. Either of the above techniques may be employed to introduce the EPSPS and /or ALS gene cassettes into tobacco or any other suitable plant tissue or cells. The general cassette consists of: (1) Drug selection/degrading marker gene such as kanamycin or any other suitable selection marker; (2) a strong promoter element such as 35sCMV

(cauliflower mosaic virus); and (3) right and left border T DNA repeats of Agrobacterium. Such a cassette could be introduced into plants either by a biolistic process or by Agrobacterium mediated gene transfer (Horsch, et al., *Nature* 227:1229-1231 (1985)). The cassette is based on pBI121 gene transfer vector (Jefferson, et al., *EMBO J.*, 6:3901-3907 (1987)). The design of the final cassette is illustrated in Figure 22.

10 In the biolistic process, the transforming DNA is coated on the surface of fine gold particles and introduced into the plant cell by a particle accelerator gun (PDS1000/He gun, Biorad, Richmond, CA). For Agrobacterium mediated gene transfer the transforming DNA cassette is introduced into the bacteria. The Agrobacterium harboring the cassette is allowed to be in contact with a disk or tissue section from tobacco or other suitable plant leaves. This facilitates the transfer of the DNA cassette to the plant nuclei. In either of the above approaches, the DNA finally gets integrated into the plant nuclei. The putative transformed cells are used for marker gene (drug) selection. The plants regenerated in presence of the selected drugs are strong transgenic candidates. After the plants are mature, the cell extracts will be taken and mixed with SDS loading dye with 1 mM DTT, boiled at 95°C for 20 5 min and loaded on a 10-20% Tris-glycine-SDS gradient gel. The separated proteins will be blotted on an Immobilon-P membrane and probed with an anti-ALS or EPSPS antibody.

PCR may then be performed to determine if the gene has integrated in a predictable fashion without rearrangement.

This technique would be useful for proteins that need specific modification for activity/folding in cytoplasmic environment. A part of the target protein gene with necessary transport signal and splicing elements will be placed in an organelle for cytoplasmic transport in the form of a precursor polypeptide.

These plants are allowed to grow in the greenhouse till they mature and the seeds will be collected. The collected seeds are then germinated and F1 plants tested for herbicide resistance. A small-scale trial may be done to see whether or not the segregation pattern of the introduced transgenes follows a Mendelian inheritance pattern. Integration into nuclear DNA would yield Mendelian inheritance, whereas integration into chloroplast DNA would yield non-Mendelian maternal inheritance.

EXAMPLE VII

Trans-splicing of a split gene, such as EPSPS/ALS or of two unrelated gene products, such as aminoglycoside-3-acetyltransferase (aadA) and soluble modified green fluorescent protein (smGFP), to give a functional hybrid protein in plant chloroplast.

The aim of these experiments is to investigate if trans-splicing is feasible in plant chloroplasts. Plant chloroplasts are

similar to bacteria with respect to their transcription and translation machinery. In Examples IV-VI, we have used the naturally occurring intein from the *dnaE* gene of the *Synechocystis* species PCC6803, which is a cyanobacterium. Cyanobacteria are photosynthetic bacteria which are similar to plant chloroplasts. Thus it should be possible for inteins to splice or trans-splice in plant chloroplasts. These proposed experiments are in two sections: Section 1, To demonstrate the trans-splicing event of two unrelated gene products *aadA* and *smGFP* in plant chloroplasts, where both genes are integrated in chloroplast genome; and Section 2, Trans-splicing in chloroplast, where the *smGFP* gene cassette is integrated into the nuclear genome and the translated protein containing a transit peptide (rubisco 3A-IN_C-*smGFP*) is imported into the chloroplast for the reaction to proceed. The chloroplast will have *aadA* gene fused to IN_n fragment. The detailed protocol is narrated below.

To demonstrate trans-splicing of two unrelated gene products, aminoglycoside-3-acetyltransferase (*aadA*) and soluble modified green fluorescent protein (*smGFP*) in chloroplast, upon transcription and translation in chloroplasts.

The plasmids are designated as p115ag and p225ag as in Example V. These plasmids will be delivered into plant organelles using a biolistic device. Tobacco or any other suitable plant tissue will be harvested aseptically from sterile greenhouse grown plants or tissue culture plant cells. Plant

tissue will be equilibrated overnight with plant growth medium and sorbitol or any other suitable osmoticum. The plant cells will be bombarded with the above plasmids coated on gold particles. After a suitable recovery time the cells will be placed on plant growth medium along with phytohormone and spectinomycin sulphate 500ug/ml. The spectinomycin resistant callus tissue will be harvested and will be placed on shoot differentiation medium. When shoots are about 2 cm length they will be dissected out and put in the rooting medium. The transgenic plant or sector of the plants will be identified by hand held UV lamp (a normal (non-transgenic) plant will fluorescent red in UV, whereas, a transgenic plant will look green). The transgene integration and copy numbers will be verified by Southern blot analysis and PCR. The transgenic sectors will be tested for trans-splicing of aadA and smGFP using anti GFP antibody. These sectors would further be used for generating a pure trans-plastomic line. The F1 plants will be tested for spectinomycin resistance.

Trans-splicing in the chloroplast. The smGFP gene cassette is integrated into the nuclear genome and the translated protein containing the transit peptide of the rubisco 3A-IN_C-smGFP is imported into the chloroplast for the reaction to proceed.

This method will enable any split protein (e.g., EPSPS or ALS) to be expressed as fused proteins with either IN_n or

IN_C either in chloroplasts or the nucleus. The nuclear-encoded component will be fused to a chloroplast transit peptide to facilitate its migration into the chloroplast after translation in the cytoplasm. A detailed method for aadA and GFP is given below. Similar methods could be followed for any other protein/split genes.

This method will require a nuclear transformation vector, such as pBI121, carrying a drug selection marker and the target gene of interest. Our experimental gene will be a three part fusion protein with rubisco transit peptide followed by IN_C and smGFP (in place of smGFP another protein/peptide such as half of EPSPS or ALS could be substituted). The transit peptide is codon optimized for tobacco (Figure 26). This fusion gene will be under the control of a strong plant promoter, 35SCMV. A diagram of such cassette is shown (Figure 23). This DNA will be introduced into the plant nucleus. The stable transgenic lines will be selected and F1 progeny will be tested for transgene integration.

Leaf sections from the above transgenic plants will be used for chloroplast DNA transformation. The chloroplast gene targeting vectors are based on p114 and p224 with spectinomycin resistance gene and a PpsbA promoter to drive the transgene. The transgenes could be the other half of the protein (that was introduced to the nuclear genome previously) with the necessary splicing elements. As a model system we would use the aadA-IN_n fusion gene for

chloroplast transformation. The transplastomic lines will be selected using both drugs (e.g., the chloroplast specific drug spectinomycin and the nuclear specific drug kanamycin). PCR and Western blot analysis will further establish pure plant lines.

For the transgenic plants the F1 generation will be tested for: (1) Mendelian inheritance pattern of the transgene/segment; (2) stability of the transgene; and (3) possible escape of the transgene through pollen.

ALS/EPSPS transgenic plants will be tested for resistance to sulphonyl urea and Roundup®.

It should be understood that the Examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be apparent to persons skilled in the art and are to be included within the spirit and purview of this Application and the scope of the appended claims.

WHAT IS CLAIMED IS:

1. A method of reconstituting a target protein in a predetermined location within an organism comprising:

5 (a) splitting DNA coding for the target protein into at least two fragments;

(b) separating the DNA fragments of step (a) to prevent transmission of the gene coding for the target protein to other organisms;

10 (c) expressing the DNA fragments of step (b) within the organism to produce the corresponding fragments of the target protein; and

(d) reconstituting the target protein from the protein fragments.

15 2. A method preventing transmission to other organisms of the gene coding for a target protein from within an organism containing said DNA coding for the target protein comprising:

20 (a) splitting DNA coding for the target protein into at least two fragments; and

(b) separating the DNA fragments of step (a) to prevent transmission of the gene coding for the target protein.

25 3. The method of claim 1 or 2, wherein the organism is selected from the group consisting of plants, animals, fungi, viruses, prokaryotes, and single-cell eukaryotes.

4. The method of claim 1 or 2, wherein the DNA coding for the target protein is split by DNA coding for one or more inteins or portions thereof.

5. The method of claim 4, wherein the DNA coding for the target protein is split by forming at least two DNA fusion fragments, wherein said DNA fusion fragments comprise a portion of the DNA coding for the target protein and a portion of DNA coding for the intein.

6. The method of claim 5, wherein one of said fusion fragments is formed by linking the C-terminal end of DNA coding for an N-terminal portion of the target protein to the N-terminal end of the DNA coding for an N-terminal portion of the intein, and another of said fusion fragments is formed by linking the N-terminal end of DNA coding for a C-terminal portion of the target protein to the C-terminal end of DNA coding for a C-terminal portion of the intein.

7. The method of claim 1 or 2, wherein the DNA coding for the target protein is split to form two or more DNA fragments by DNA coding for one or more affinity domains.

8. The method of claim 7, wherein the affinity domain is selected from the group consisting of inteins or intein fragments, leucine zipper and c-Jun/c-Fos.

9. The method of claim 1 or 2, wherein the DNA fragments coding for the target protein are separated by compartmentalizing each DNA fragment into different compartments selected from a group consisting of the nucleus, a membrane bound organelle, a plasmid, a virus, a cosmid, and an artificial chromosome.

10. The method of claim 9, in which at least one of the DNA fragments coding for the target protein is fused to a DNA sequence encoding transit peptides such that the protein products of the DNA fragments are transported into a single compartment where functional reconstitution can occur.

11. The method of claim 10, in which one of the DNA fragments coding for a portion of the target protein is compartmentalized in the nucleus, being fused to a DNA sequence encoding a transit peptide for transport into chloroplasts, and the other DNA fragment coding for another portion of the target protein is compartmentalized in the chloroplasts.

12. The method of claim 1 or 2, wherein the DNA fragments coding for the target protein are separated by inserting each of the fragments into different portions of a DNA molecule wherein the DNA molecule is selected from the group consisting of DNA from the nucleus, a membrane bound organelle, DNA from a plasmid, DNA from a cosmid, DNA from a virus and DNA from an artificial chromosome.

13. The method of claim 12, wherein at least one of the DNA molecules is naturally inherited.

5 14. The method of claim 12, wherein at least one of the DNA molecules resides in the chloroplasts.

15. The method of claim 12, wherein at least one of the DNA molecules resides in the mitochondria.

10

16. The method of claim 4, wherein reconstitution of the target protein fragments comprises intein-mediated splicing.

15

17. The method of claim 4, wherein reconstitution of the target protein fragments comprises intein-mediated protein complementation.

20

18. The method of claim 1, wherein reconstitution of the target protein fragments comprises protein complementation.

19. The method of claim 18, wherein protein complementation occurs in the presence of an affinity domain.

25

20. The method of claim 18, wherein protein complementation occurs in the absence of an affinity domain.

21. The method of claim 1 or 2, wherein splitting of the DNA coding for the target protein comprises:

- (a) determining one or more potential split site regions of the target protein; and
- (b) splitting the DNA coding for the target protein at the potential split site region.

5

22. The method of claim 21, wherein the potential split site region of the target protein is determined by analyzing primary amino acid sequence of the target protein for non-conserved regions.

10

23. The method of claim 21, wherein the potential split site region is determined by linker tolerance of linker insertion within the target protein.

15

24. The method of claim 21, wherein the potential split site region is determined by analyzing the structure of the target protein for the presence of flexible loops.

20

25. The method of claim 21, wherein the potential split site region is determined by analyzing the structure of the target protein for the presence of amino acid sequence between folding domains of the target protein.

25

26. An isolated DNA fragment comprising a DNA split site in an EPSPS gene.

27. The isolated DNA fragment of claim 26, wherein the DNA fragment is selected from the group consisting of the DNA

encoding for amino acids 1-235 or portions thereof, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38 and SEQ ID NO:39.

5

28. An isolated DNA fragment comprising a DNA split site in an *E. coli* ALS gene.

10

29. The isolated DNA fragment of claim 28, wherein the DNA fragment is selected from the group consisting of SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12 and SEQ ID NO:13.

15

30. An isolated DNA fragment comprising a DNA split site in a maize ALS gene.


31. The isolated DNA fragment of claim 30, wherein the DNA fragment is selected from the group consistin of SEQ ID NO:17, SEQ ID NO:17, SEQ ID NO:18 and SEQ ID NO:19.

20

32. The isolated DNA fragments of claim 26, 28, or 30, wherein said DNA fragment is fused to DNA coding for an intein or portion thereof.

1/33

FIG. 1A



2/33

FIG. 1B





FIG. 2A



3/33

FIG. 2B






FIG. 3



4/33

FIG. 4A



5/33

FIG. 4B





FIG. 5



6/33

FIG. 6

327 Y A V D K A D L L L A L G V R F D D R V T G K L E A F A S R Maize ALS
 356 Y A V D S S D L L L A F G V R F D D R V T G K L E A F A S R Tobacco ALSI
 353 Y A V D S S D L L L A F G V R F D D R V T G K L E A F A S R Tobacco ALSII
 268 M T M H N A D V I F A V G V R F D D R T T N N L A K Y C P N E. Coli ALSIII
 258 F A V Q E C D L L I A V G A R F D D R V T G K L N T S A P H E. Coli ALSII

357 A K I V H V D I D P A E I G K N K Q P H V S I C A D V K L A Maize ALS
 386 A K I V H I D I D S A E I G K N K Q P H V S I C A D I K L A Tobacco ALSI
 383 A K I V H I D I D S A E I G K N K Q P H V S I C A D I K L A Tobacco ALSII
 298 A T V L H I D I D P T S I S K T V T A D I P I V G D A R Q V E. Coli ALSIII
 288 A S V I H M D I D P A E M N K L R Q A H V A L Q G D L N A L E. Coli ALSII

*

387 L Q C M N A L L E G S T S K K S F D - F G S W N D E L D O Q Maize ALS
 416 L Q G L N S I L E S K E G K I K L D - F S A W R Q E I T E Q tobacco ALSI
 413 L O G L N S I L E S K E G K I K L D - F S A W R Q F I T V O tobacco ALSII
 328 L E Q M L E L L S Q E S A H Q P L D E I R D W W Q Q I E Q W E. Coli ALSIII
 318 L P A L Q Q P L N Q C D - - - - - W Q Q H C A Q L E. Coli ALSII


416 K R E F P L G Y K T S N E E I Q P Q Y A I Q V L D E I T K G Maize ALS
 445 K V K H P L N F K T F G D A I P P Q Y A I Q V L D E I T N G tobacco ALSI
 442 K V K Y P L N F K T F G D A I P P Q Y A I Q V L D E I T N G tobacco ALSII
 358 R A R Q C L K Y D T H S E K I K P Q A V I E T L W R I T K G E. Coli ALSIII
 338 R D E H S W R Y D H P G D A I Y A P L L L K Q L S D R K P A E. Coli ALSII

446 E A I I G T G V G Q H Q M W A A Q Y Y T Y K R P R Q W L S S Maize ALS
 475 N A I I S T G V G Q H Q M W A A Q Y Y K Y R K P R Q W L T S tobacco ALSI
 472 S A I I S T G V G Q H Q M W A A Q Y Y K Y R K P R Q W L T S tobacco ALSII
 388 D A Y V T S D V G Q H Q M F A A L Y Y P F D K P R R W I N S E. Coli ALSIII
 368 D C V V T T D V G Q H Q M W A A Q H I A H T R P E N F I T S E. Coli ALSII

476 A G I G A M G F G L P A A A G A S V A N P G V T V V D I D G Maize ALS
 505 C G I G A M G F G L P A A I G A A V G R P D E V V V D I D G tobacco ALSI
 502 G G I G A M G F G L P A A I G A A V G R P D E V V V D I D G tobacco ALSII
 418 G G I G T M G F G L P A A L C V K M A L P E F T V V C V T C E. Coli ALSIII
 398 S G I G T M G F G L P A A V G A Q V A R P N D T V V C I S G E. Coli ALSII

7/33

FIG. 7



8/33

FIG. 8A





FIG. 8B



FIG. 8C





9/33

FIG. 9A



10/33

FIG. 9B



11/33

FIG. 10A






FIG. 10B




12/33

FIG. 11



13/33

FIG. 12



14/33

FIG. 13A






FIG. 13B



15/33

FIG. 14



16/33

FIG. 15-1

EPSPS Insertion Site	Amino acid sequence inserted	Clone
Q7/P8	CLNIO	pCE-5aa 129
A10/R11	VFKHA	pCE-5aa 47
P35/C36	LFKQP	pCE-5aa 7
D48/D49	CLNSD	pCE-5aa 50
S67/A68	CLNIS	pCE-5aa 8
D69/R70	CLNTD	pCE-5aa 44
R70/T71	CLNNR	pCE-5aa 10
C73/D74	CLNSC	pCE-5aa 32
D74/I75	CLNSD	pCE-5aa 5
L82/R83	CLNTL	pCE-5aa 3
P85/G86	VFKQP	pCE-5aa 12
M121/K122	CLNSM	pCE-5aa 42
Y148/P149	CLNNY	pCE-5aa 37
L182/A183	CLNTL	pCE-5aa 22
A183/P184	CLNMA	pCE-5aa 11
K185/D186	VFKHK	pCE-5aa 112
K185/D186	CLNTK	pCE-5aa 212
D186/T187	CLNKD	pCE-5aa 33
I188/I189	MFKOI	pCE-5aa 151
I189/R190	CLNII	pCE-5aa 114
E194/L195	LFKHE	pCE-5aa 227
F211/G212	VFKHF	pCE-5aa 162
V213/E214	CLNSV	pCE-5aa 1
I215/A216	VFKOI	pCE-5aa 2
A216/N217	MFKOA	pCE-5aa 208
H218/H219	LFKHH	pCE-5aa 28
Q221/Q222	LFKHQ	pCE-5aa 4
V225/K226	MFKHV	pCE-5aa 203
K226/G227	VFKOK	pCE-5aa 25
Q230/Y231	LFKQO	pCE-5aa 102
S233/P234	LFKHS	pCE-5aa 40
G235/R236	CLNTG	pCE-5aa 35
R267/K268	CLNSR	pCE-5aa 23
L238/V239	VFKHL	pCE-5aa 154

17/33

FIG. 15-2


EPSPS Insertion Site	Amino acid sequence inserted	Clone
I311/P312	CLNNI	pCE-5aa 29
Q375/H376	LFKHO	pCE-5aa 15
Q375/H376	CLNIO	pCE-5aa 223
H376/A377	CLNKH	pCE-5aa 38
Y382/N383	MFKOY	pCE-5aa 31
E418/Q419	LFKHE	pCE-5aa 36
Q419/L420	CLNKO	pCE-5aa 46
S424/T425	CLNMS	pCE-5aa 9

FIG. 16

EPSPS Insertion Site	Amino acid sequence inserted	Clone
L31/A32	LCLNILA	pCE-5aa 21d
N55/A56	NCLNINA	pCE-5aa 4d
L57/S58	LMFKHLS	pCE-5aa 217
T71/R72	TLFKHTR	pCE-5aa 24d
K122/E123	KVFKOKE	pCE-5aa 126
H128/L129	HLVFKHL	pCE-5aa 142
L176/L177	LCLNTLL	pCE-5aa 122
L238/V239	LCLNNLV	pCE-5aa 205
E240/G241	EVFKHEG	pCE-5aa 171
K256/G257	KVFKQKG	pCE-5aa 140
T286/I287	TCLNTTI	pCE-5aa 180
M328/N329	MCLNNMN	pCE-5aa 115
L331/R332	LLFKQLR	pCE-5aa 124
R344/L345	RCLNNRL	pCE-5aa 107
M348/A349	MVFKQMA	pCE-5aa 3d
A349/T350	AMFKQAT	pCE-5aa 110
L404/D405	LVFKHLD	pCE-5aa 199
K411/T412	KMFQOKT	pCE-5aa 5d
Y416/F417	YCLNNYF	pCE-5aa 163


18/33

FIG. 17



19/33

FIG. 18



EXPRESSION PLASMID pAGR3: 5910 bp.
PROMOTER AND CLONING SITE MAP:

lac operator

1 GAATTGTGAG CGCTACAAT TCTAGGATGT TAATTGCGCC GACATCATAA

-35 region

51 CGGTTCTGGC AAATATTCTG AAATGAGCTG TTGACAATTA ATCATCGGCT

-10 region

lac operator

rbs

101 CGTATAATGT GTGGAATTGT GAGCGGATAA CAATTCACA CAGGAAACAG

start

151 ACCATGGTGA ATTCTAGAGC TCGAGGATCC GCGGTACCCG GGCATGCATT
NcoI EcoRI XbaI SacI XhoI BamHI SacII KpnI SmaI BstBI

201 CGAAGCTTCC TTAAGCGGCC GTCGACCGAT GCCCTTGAGA GCCTTCAACC
HindIII AflIII EagI SalI

20/33
FIG. 19A




FIG. 19B




FIG. 19C





FIG. 20



22/33

FIG. 21A




FIG. 21B

	Top10 E. coli	pAGRECsmGFP	p115ag4	p115ag11	p225ag3	p225ag12
Amp	-	+	+	+	+	+
Amp+Spen	-	-	+	+	+	+

FIG. 21C




FIG. 22





FIG. 23



25/33

FIG. 24

GAATAGATCTACATACACCTTGGTTGACACGAGTATATAAGTCATGTT
ATACTGTTGAATAACAAGCCTTCCATTTTCTATTTTGATTGTAGAAA
ACTAGTGTGCTTGGGAGTCCCTGATGATTAAATAAACCAAGATTTTAC
CTTAATTAAG

FIG. 25

GATCCTGGCCTAGTCTATAGGAGGTTTTGAAAAGAAAGGAGCAATAAT
CATTTTCTTGTCTATCAAGAGGGTGCTATTGCTCCTTTCTTTTTTC
TTTTTATTTATTTACTAGTATTTTACTTACATAGACTTTTTTGTTTAC
GTATTCT

FIG. 26

catATGGCgTccATGATcTCCTCgTCcGCgGTGACcACgGTCAGCCGcG
CgTCcACGGTGCAgTCGGCCGCGGTG6CcCCgTTCGGCGGCCTCAAgTC
CATGACcGGcTTCCcGgTcAAGAAGGTCAACACgGACATcACgTCCATc
ACgAGCAAcGGcGGcAGgGTgAAGTGCATGcgaagagc

FIG. 27-1

GTAACTACGTCAGGTGGCACTTTTCGGGAAATGTGCCGGAACCC
CTATTTGTTATTTTCTAAATACATTCAAATATGTATCCGCTCATG
AGACAATAACCCTGATAAATGCTTCAATAATTGAAAAAGGAAGAG
TATGAGTATTCAACATTTCGCTGTCGCCCTTATTCCTTTTTCGCG
CATTTTGCCTTCTGTTTTGCTCACCAGAAACGCTGGTGAAGTA
AAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAAT
GGATCTCAACAGCGGTAAGATCCTTGAGAGTTTTGCCCCGAAGAAC
GTTCTCAATGATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGTA
TTATCCCGTGTGACGCCGGCAAGAGCAACTCGGTGCCGCATACA
CTATTCTCAGAAATGACTTGGTTGAGTACTACCAAGTCACAGAAAAGC
ATCTTACGATGGCATGACAGTAAGAGAATTATGAGTGTGCCATA
ACCATGAGTGATAACACTGCCGCCAACTTACTTGAACAAGATCGG
AGGACCGAAGGAGCTAACCGCTTTTTCACAACATGGGGGATCATG
TAACTCGCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACCA
AACGACGAGCGTGACACCACGATGCCGTGAGCAATGGCAACAACGTT
GCGCAAACTATTAACTGGCGAACTACTTACTTAGCTTCCGGCAAC
AATTAATAGACTGGATGGAGGCGGATAAAGTTGCAGGACCACTTCTG
CGCTCGGCCCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGC
CGGTGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATG
GTAAGCCCTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCA
ACTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTCACT
GATTAAGCATTGGTAAGTGTGAGCAAGTTTACTCATATATACTTT
AGATTGATTTACCCCGTTGATAATCAGAAAAGCCCCAAAACAGGA
AGATTGTATAAGCAAATATTTAAATTGTAACGTTAATATTTTGTTA
AAATTCGCGTTAAATTTTGTAAATCAGCTCATTTTAAACCAATA
GGCCGAAATCGGCAAAATCCCTTATAAATCAAAGAATAGCCCGAGA
TAGGGTTGAGTGTGTTCCAGTTTGGAAACAAGAGTCCACTATTAAAG
AACGTGCACTCCAACGTCAAAGGGCGAAAACCGTCTATCAGGGCGA
TGGCCCACTACGTGAACCATCACCAAAATCAAGTTTTTGGGGTCGA
GGTGCCGTAAGCACTAAATCGGAACCTAAAGGGAGCCCCGATTT
AGAGCTTGACGGGGAAGCGAACGTGGCGAGAAAGGAAGGGAAGAAA
GCGAAAGGAGCGGGCGCTAGGGCGCTGGCAAGTGTAGCGGTACGCT
GCGCGTAACCAACACCCCGCGCTTAATGCCCGCTACAGGGCG
CGTAAAGGATCTAGGTGAAGATCCTTTTGTAAATCTCATGACCAA
AATCCCTTAACGTGAGTTTTCGTTCCACTGAGCGTCAGACCCGATG
AAAAGATCAAAGGATCTTCTGAGATCCTTTTCTGCGCGTAATC
TGCTGCTTGCAAAACAAAAAACCCGCTACCAAGCGGTGTTGTTT
GCCGGATCAAGAGCTACCAACTCTTTTCCGAAGGTAAGTGGCTTCA
GCAGAGCGCAGATACCAATACTGTTCTTCTAGTGTAGCGTAGTTA
GGCCACCACTTCAAGAACTCTGTAGCAGCGCTACATACCTGCTCT
GCTAATCCTGTTAC

27/33

FIG. 27-2

CAGTGGCTGCTGCCAGTGGCGATAAGTCGTGCTTACCGGGTTGGA
CTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTCGGGCTGAACG
GGGGGTTCTGTGCACACAGCCAGCTTGGAGCGAACGACCTACACCG
AACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGCTTCC
CGAAGGGAGAAAGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGA
ACAGGAGAGCGCACGAGGGAGCTTCCAGGGGAAACGCCTGGTATC
TTTATAGTCTGTGCGGTTTCGCCACCTCTGACTTGAGCGTCGATT
TTTGTGATGCTCGTCAGGGGGCGGAGCCTATGAAAAACGCCAGC
AACCGGGCTTTTTACGGTTCTTGGCCTTTTGCTGGCCTTTTGCTC
ACATGTAATGTGAGTTAGCTCACTCATTAGGCACCCAGGCTTTAC
ACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGGATA
ACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTA
CGTAATACGACTCACTAGTGGGCAGATCTTGAATGCATCGCGCGC
TTGACGATATAGCAATTTTGCTTGGATTTATCAGTCGAAGCAGGAG
ACAATATACCTTGATATTCTCGATCATTCTTTGATTCAAAGCATCG
TTCCATCTCAATTGAAAAAGCAAATAACGTTTCAAGAACAAATCTA
GTTCTGCTTCCGTGTGCTTTTGATTGTTTTTCTTTTACCCTT
CTTTGTGCTGATTCCGCGTAATCTTTTTAAGAGCGTTTGATGT
TTTGAGAGAACAGGGCCAGATTTCTTTGTTTTCTATATCTGATC
CACGCTCTTTTTCTCCTTGACTTGCGGGTCTTTTGCTTCTTGAAT
TCGATTCTTTATTTTTTTTATTGATCGTAGAAAAAGTTTTGTTTT
TGGTTTTTATTGATGTTTTTATTGACTAACATTTTCATTGAT
TCAAATTTAAAGAAATTTGCTTGGTATAATCCACGGTTTTAT
TTTATATACATTATAAAGTGGTACAAATCTGGGAAGAACCAAAAT
TCCAGATTCAATATGGGACGATTAATATTTTTTCATTATTCCCA
TCCAATCAAAAAGGCTTTTTTCGAATTTTTTGATTGTTTTCTGG
ATTTTGATGAATCGTAAGATAAAAAAGCCTTTTTTATCAATTTTA
TCAATTATTTGATAATTATTAATACCAATTTTAGTATTTGGATTAC
TGTTGGTATCGATCTTAACCCAGGCCTCAATATCTTCTTTTGTCT
AAGAGAAAAATGGATAATTTTCAATCAAAATATTTTCTATCGAGA
TTTCTTTCTATATATAGAATATTGCCTTTTCTTAGATAATTATTGA
TATGAAGATTGCCGAGCATATCAAAAAGGTTGTGTTGGACGTGT
GGAATTAGAAGAAATTTGAGGTTCTTATTTACTTGAAAGGGTAAT
CTAGAAATAAAGAGTCATTTTTTTTTCATAATTAATCGATTTAT
ATGCTAAAAGATCATATCTATAACATTTTTGAAAATTATCTTTTG
GTTTGCTAATGAATAGAGCTCAGAAATCATTTTCTTTTTTGTAATGA
ATTAATTGGTCTTTTTCATATGAATTCATTTGTTTAAATTTTCGAT
TTTGAGCCATACAACCTTGATTAACCCTATTTGCCATTTTGTGG
CATTAAATCTAGACCATCTAATCTGAGATAAATCGTACGagaatact
caatCATGAATAAATGCAAGAAAATAACCTCTCCTTTCTTTCTAT
AATGTAACAAAAAGTCTATGTAAGTAAATACTAGTAAATAAAT
AAAAAGAAAAAAGAAAGGAGCAATAGCACCCCTCTTGATAGAACAA
GAAATGATTAT

28/33

FIG. 27-3

TGCTCCTTTCTTTTCAAACCTCCTATAGACTAGGCCAGGATCCTCGA
GcttaattaaGGTAAATCTTG6TTTATTTAATCATCAGGGACTCCCA
AGCACACTAGTTTTCTACAAATCAAATAGAAAATAGAAAATGGAAGG
CTTTTTATTCAACAGTATAACATGACTTATATACTCGTGTCAACCAAG
GTGTATGTAGATCtattcCTGCAGGATATCTGGATCCACGAAGCTTCC
CATGGGAATAGATCTACATACACCTTGGTTGACACGAGTATATAAGTC
ATGTTATACTGTTGAATAAAAAGCCTTCCATTTTCTATTTTGATTGT
AGAAAAGTGTGCTTGGGAGTCCCTGATGATTAATAAACCAAGAT
TTTACCGTTTAAACACCGGTGATCCTGGCCTAGTCTATAGGAGGTTTT
GAAAAGAAAGGAGCAATAATCATTTTCTTGTTCTATCAAGAGGGT6CT
ATTGCTCCTTTCTTTTTCTTTTATTTATTTACTAGTATTTACTT
ACATAGACTTTTTGTTTACATTATAGAAAAAGAGGAGGTTATTT
TCTTGCAATTATTATGATTGAGTATTCTcctaggCGATTGATAATG
CCGCTTAACCAAGTTTTCCATTGATTGATTCTATAACTCTGAAGTTT
CTTATGTTTAAATTCAGAAATGAAATATTCTAGTGTTCGAAAATAGTC
CTTTATTTAGTCTTAAGGAAAAAGACGTTCTGTTATATTGAAGAAC
AGATCTTAATTTAGACAAATTAATAACTTGGGGTTGTGATAATTTGTA
AAATACATATGCTTGTGATAAGTAGGATAAATCAAAAAAATATGTGA
ATTTTTCTTACTAATATTATAAAGTGACTTTTTTATAGTCGAAATAAA
GTGAATTTTTTTTGATTATTATTTTTCTTGATTATTTTCAATTATT
GGAAATGTATTTATCAATCAATTTGTTTGTGATTCAAGAAAGAGTTG
TGTATTAATCTGGAATATTAATGATAGATAAAATAGATCGATGTA
TAATCTTTGAATGAATAATTTAGAAAATGGAATTTCCATATTAA
TCGAGTATTTCTTCTTTTAAATATTGGAATCTTTTTGCGGATTC
GAATTTTTAAATATTATTGTTTTATTAGGACTAATGTCTATTTCTG6
AGTTACTTTCTTTTCTCTTTTGAATCTTTCTATTTGATTTTGTGAT
TGTACTTGTCTATCAGTCAAATCCTTCATTTTGTCTTCTATCAGTGA
AGAATTTGGCCAATTTCCAGATTCAATTTGACTAAATGATTGTTAAT
TATCTGATTACTCATTAGAGAATCTTTTCTTTTTCTGTTTCATTGGA
TTCATCTATTTCTTTGAGTCTAAATAATACAATTGGATTTACTTTTGA
AAGTCTTTTTCTTTTTTTTATAAATAGACTACTTTTGATAAGCCA
TTTTTTGGTTTCTTTGAAATCTTCGAAATAATTTATTTTCTTTT
GAAAACCTTTAGAGTTATAAAATATTTCTTTTGAATTTTCAATTTT
TTTTTCGAGTTCCTTAAAAATGGGCTCAAAAAAGAGGGGCTTTTCG
GGGAGAACCAGGGAAGTTCAGCTTCCATTCCCAAACTGTTAAAAA
ACAAAAATCATCTTTTGTTTTTCTTTTTCATTAGCTCTCCACGGGA
GGAGTACAGTTAGATATATGCCAAGGTTTCAGACAAAAAGGAAATAA
TATTTTGATCTGAATGCCATCTTCAACCAATTTTTGGAATTTCTGT
TTCTGATAATTGAACACCATTATAAGTACATTTAATATGCATTTCTCT
ATTCCATTCTGCAATCTTCAGACCATTGAGGAAGTTGCAAGACTAA
CATACGCCGAGATTTTGGCTATTATCAATGAAGGTAATACAATATA
TTTTCGAAGAATTG

29/33

FIG. 27-4

ATTGAGTTATTAACATGTAACCTCTTATTATTTGCGCAAAAGGAATGGT
ATCCCAGGCTTCTGCTATCTCTATCCGTGCTTTTTCCTTTCTTTTGTTT
TCCCCTTTTTGTCCTTTTCCTTTTCTCTCTCTTTTGTGTTCTT
CTCTAGACTCTAGAATCTTGAATTCGGTACCTCTAGTCAAGGCCTTAA
GTGAGTCGTATTACGGAAGTGGCCGTCGTTTACAACGTCGTGACTGGGA
AAACCCTGGCGTTACCCAACCTTAATCGCCTTGACGACATCCCCCTTC
GCCAGCTGGCGTAATAGCGAAGAGGCCCGCACCAGTCGCCCTTCCCAAC
AGTTGCGCAGCCTGAATGGCGAATGGCGCTTCGCTTGGTAATAAGGCC
GCTTCGGCGGGCTTTTTTTT

30/33

FIG. 28-1

GTAACTACGTACGGTGGCACTTTTCGGGAAATGTGCGGGAACC
CCTATTTGTTATTTTCTAAATACATTCAAATATGTATCCGCTCA
TGAGACAATAACCGTGATAAATGCTTCAATAATATTGAAAAGGAA
GAGTATGAGTATTCAACATTTCCGTGTCGCCCTATTCCCTTTTT
GCGGCATTTTGCTTCTGTTTTGCTCACCAGAAACGCTGGTGA
AAGTAAAAGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACAT
CGAAGTGGATCTCAACAGCGGTAAAGATCCTTGAGAGTTTTGCCCC
GAAGAACGTTCTCCAATGATGAGCACTTTAAAGTTCTGCTATGTG
GCGCGGTATTATCCCGTGTGACGCCGGGCAAGAGCAACTCGGTG
CCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTACCAAGTC
ACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCA
GTGCTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCT
GACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTCACAAC
ATGGGGGATCATGTAAGTACGCTTGATCGTTGGGAACCGGAGCTGA
ATGAAGCCATACCAAACGACGAGCGTGACACCAGATGCTGTAGC
AATGGCAACAACGTTGCGCAAACTATTAAGTGGCGAACTACTTACT
CTAGCTTCCCGGCAACAATTAATAGACTGGATGGAGGCGGATAAG
TTGAGGACCACTTCTGCGCTCGGCCCTTCGGCTGGCTGGTTTAT
TGCTGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTATCATT
GCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTATCT
ACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGAT
CGCTGAGATAGGTGCTCACTGATTAAGCATTGGTAAGTGTGAGAC
CAAGTTTACTCATATATACTTTAGATTGATTTACCCCGGTTGATAA
TCAGAAAAGCCCCAAAACAGGAAGATTGTATAAGCAATATTTAA
ATTGTAAACGTTAATATTTTGTAAAATTCGCGTTAAATTTTGT
AAATCAGCTCATTTTTTAACCAATAGGCCGAAATCGGCAAAATCCC
TTATAAATCAAAAGAATAGCCCGAGATAGGGTTGAGTGTGTTCCA
GTTTGAACAAGAGTCCACTATTAAGAACGTTGACTCCAACGTCA
AAGGGCGAAAAACCGTCTATCAGGGCGATGGCCCACTACGTGAACC
ATCACCCAAATCAAGTTTTTGGGGTCGAGGTGCCGTAAAGCACTA
AATCGGAACCCCTAAAGGGAGCCCCGATTAGAGCTTGACGGGGAA
AGCGAACGTGGCGAGAAAGGAAGGAAGAAAGCGAAAGGAGCGGGC
GCTAGGGCGCTGGCAAGTGTAGCGGTACGCTGCGCGTAACACCA
CACCAGCGCGCTTAATGCGCCGCTACAGGGCGCGTAAAGGATCT
AGGTGAAGATCCTTTTGTAAATCTCATGACCAAAATCCCTTAACG
TGAGTTTTCTGTTCCACTGAGCGTCAGACCCGTAGAAAAGATCAA
GGATCTTCTTGAGATCCTTTTTTCTGCGCGTAATCTGCTGCTTGC
AAACAAAAAACACCGCTACCAGCGGTGGTTTGTGTTGCGGATCA
AGAGCTACCAACTCTTTTTCCGAAGGTAAGTGGCTCAGCAGAGCG
CAGATACCAAACTGTTCTTCTAGTGTAGCCGTAGTTAGGCCACC
ACTTCAAGAACTCTGTAGCACCCTACATACCTCGCTCTGCTAAT
CCTGTTAC

31/33

FIG. 28-2

CAGTGGCTGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGTTGGA
CTCAAGACGATAGTTACCGGATAAGGCGCAGCGGTGGGGTGAACG
GGGGGTTCTGTCACACAGCCAGCTTGGAGCGAACGACCTACACCG
AACTGAGATACCTACAGCGTGAGCTATGAGAAAGCCACGCTTCC
CGAAGGGAGAAAGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGA
ACAGGAGAGCGCACGAGGGAGCTTCCAGGGGAAACGCCCTGGTATC
TTTATAGTCCTGTGCGGTTTCGCCACCTCTGACTTGAGCGTCGATT
TTTGTGATGCTCGTCAGGGGGCGGAGCCTATGGAAAAACGCCAGC
AACGCGGCCTTTTACGGTTCTTGGCTTTTGTGGCTTTTGTCTC
ACATGTAATGTGAGTTAGCTCACTCATTAGGCACCCAGGCTTTAC
ACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGGATA
ACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTA
CGTAATACGACTCACTAGTGGGCAGATCTTGAATGCATCGCGCGC
AATTCACCGCCGTATGGCTGACCGGCGATTACTAGCGATTCCGGCT
TCATGCAGGCGAGTTGCAGCCTGCAATCCGAACAGGACGGGTTT
TTGGGGTTAGCTCACCTCGCGGGATCGGACCCTTTGTCCCGCC
ATTGTAGCACGTGTGTGCGCCAGGGCATAAGGGGCTGATGACTTG
ACGTATCCTCACCTTCTCCGGCTTATCACCGGCAGTCTGTTGAG
GGTTCCAAACTCAACGATGGCAACTAAACACGAGGGTTGCGCTCGT
TGCAGGACTTAACCAACACCTTACGGCACGAGCTGACGACAGCCA
TGCACCACCTGTGTCCGCGTTCCCGAAGGCACCCCTCTCTTTCAAG
AGGATTGCGGGCATGTCAAGCCCTGGTAAGGTTCTTGGCTTTCAT
CGAATTAACACATGCTCCACCGCTTGTGCGGGCCCCGTCAATT
CCTTTGAGTTTCACTTGTGCAACGTAATCCCGGGGATGACTT
AACGCGTTAGCTACAGCACTGCACGGGTCGATACGCACAGCGCCTA
GTATCCATCGTTTACGGCTAGGACTACTGGGGTATCTAATCCCATT
CGCTCCCCTAGCTTTCGTCTCTCAGTGTGAGTGTGCGCCAGCAGA
GTGCTTTCGCGGTTGGTGTCTTCCGATCTCTACGCATTTACCG
CTCCACCGGAAATCCCTCTGCCCCTACCGTACTCCAGCTTGGTAG
TTTCCACCGCCTGTCCAGGGTTGAGCCCTGGGATTGACGCGGGAC
TTAAAAAGCCACCTACAGACGCTTACGCCCAATCATTCCGGATAA
CGCTTGCATCCTCTGTATTACCGCGGCTGCTGGCACAGAGTTAGCC
GATGCTTATTTCCAGATACCGTCATTGCTTCTTCCGGGAAAAG
AAGTTCACGACCCGTGGGCTTCTACCTCCAGCGGCATTGCTCCG
TCAGCTTTCGCCCATTGCGGAAAATTTCCCACTGCTGCCTCCGTA
GGAGTCTGGGCGGTGTCTCAGTCCAGTGTGGCTGATCATCCTCTC
GGACCAGCTACTGATCATCGCCTTGGTAAGCTATTGCCTACCAAC
TAGCTAATCAGACGCGAGCCCTCCTCGGGCGGATTCTCTTTTG
CTCCTCAGCCTACGGGTATTAGCAGCCGTTTCCAGCTGTTGTTCC
CCTCCCAAGGGCAGGTTCTACGCGTTACTACCCGTCGCGCACTG
GAAACACCACTTCCCGTCCGACTTGCATGTGTTAAGC

32/33

FIG. 28-3

ATGCCGCCAGCGTTTCATCCTGAGCCAGGATCGAACTCTCCATGAGAT
TCATAGTTGCATTACTTATAGCTTCCTTGTTCGTAGACAAAGCGGAT
TCGGAATTGTCTTTTCATTCCAAGGCATAACTTGTATCCATGCGCTTC
ATATTCGCCCAGGAGTTCGCTCCAGAAATATAGCCATCCCTGCCCCC
TCACGTCAATCCACGAGCCTCTTATCCATTCTCATTGAACGACGGC
GGGGGAGCAAATCCAAC TAGAAAACTCACATTGGGCTTAGGGATAA
TCAGGCTCGAACTGATGACTTCACCACGTCAAGGTGACACTCTACC
GCTGAGTTATATCCCTTCCCCGCCCATCGAGAAATAGAACTGACTA
ATCCTAAGTCAAAGGCGTACGagaatactcaatCATGAATAAATGCA
AGAAAATAACCTCTCCTTCTTTTCTATAATGTAACAAAAAAGTCT
ATGTAAGTAAAACTAGTAAATAAATAAAAAAGAAAAAAGAAAGGA
GCAATAGCACCCCTCTTGATAGAACAGAAAAATGATTATGCTCCTTT
CTTTTCAAACCTCCTATAGACTAGGCCAGGATCCTCGAGcttaatt
aaGGTAAATCTTGGTTTATTTAATCATCAGGGACTCCCAAGCACAC
TAGTTTTCTACAAATCAAATAGAAAATAGAAAATGGAAGGCTTTTT
ATTCAACAGTATAACATGACTTATATACTCGTGTCAACCAAGGTGTA
TGATAGTCtattcCTGCAGGATATCTGGATCCACGAAGCTTCCCATG
GGAATAGATCTACATACACCTTGTTGACACGAGTATATAAGTCATG
TTACTGTTGAATAAAAAGCCTTCCATTTTCTATTTTGATTGTAG
AAAAGTAGTGCTTGGGAGTCCCTGATGATTAATAAACCAAGATT
TTACCGTTTAAACACCGGTGATCCTGGCCTAGTCTATAGGAGTTTT
GAAAAGAAAGGAGCAATAATCATTTTCTTGTCTATCAAGAGGGTGC
TATTGCTCCTTTCTTTTTTCTTTTTATTTACTAGTATTTTAC
TTACATAGACTTTTTTGTTTACATTATAGAAAAGAAGGAGAGGTTA
TTTTCTGCATTTATTCATGATTGAGTATTCTcctaggGTCGAGAAA
CTCAACGCCACTATTCTTGAACAACCTGGAGCCGGGCTTCTTTTCG
CACTATTACGGATATGAAAATAATGGTCAAAATCGGATTCAATTGTC
AACTGCCCCATCGGAAATAGGATTGACTACCGATTCCGAAGGAAC
GGAGTTACATCTCTTTTCCATTCAAGAGTCTTATGCGTTTCCACGC
CCCTTTGAGACCCCGAAAAATGGACAAATTCCTTTTCTTAGGAACAC
ATACAAGATTGCTCACTACAAAAAGGATAATGGTAACCTACCATT
ACTACTTCATTTATGAATTTATAGTAATAGAAATACATGCTTACC
GAGACAGAATTTGGAACCTTGCTATCCTCTTGCCTAGCAGGCAAGAT
TTACCTCCGTGGAAAGGATGATTCAATTCGGATCGACATGAGAGTCCA
ACTACATTGCCAGAATCCATGTTGTATTTTGAAGAGGTTGACCTC
CTTGCTTCTCTCATGGTACACTCCTTCCCCGCCGAGCCCTTTTCT
CCTCGGTCCACAGAGACAAATGTAGGACTGGTGCCAACAATTCATC
AGACTCACTAAGTCGGGATCACTAACTAATACTAATCTAATAATA
GTCTAATATATCTAATAATAAGAAAATACTAATAATAAGAAAAGA
ACTGTCTTTTCTGTATACTTTCCCGGTTCCGTTGCTACCGCGGGCT
TTACGCAATCGATCGGATTAGATAGATATCCCTTCAACATAGGTCAT
CGA

FIG. 28-4

AAGGATCTCGGAGACCCACCAAAGTACGAAAGCCAGGATCTTTCAG
AAAACGGATTCTATTCAAAGAGTGCATAACCGCATGGATAAGCTC
ACACTAACCCGTCAATTTGGGATCCAAATTCGAGATTTTCCTTGGG
AGGTATCGGGAAGGATTTGGAATGGAATAATATCGATTCATACAGA
AGAAAAGGTTCTCTATTGATTCAAACACTGTACCTAACCTATGGGA
TAGGGATCGAGGAAGGGGAAAAACCGAAGATTTACATGGTACTTT
TATCAATCTGATTTATTTCTGACCTTTCTGTTCAATGAGAAAATGGG
TCAAATCTACAGGATCAAACCTATGGGACTTAAGGAATGATATAA
AAAAAGAGAGGGGAAAAATTCATATTAATAAATATGAAGTAGAA
GAACCCAGATTCCAAATGAACAAATCAAACCTGAAAAGGATCTTC
CTTATTCTTGAAGAATGAGGGCAAAGGGATTGATCAAGAAAGATC
TTTTGTTCTTCTATATATAAGATCGTGATGGTACCCTCTAGTCAA
GGCCTTAAGTGAGTCGTATTACGGACTGGCCGTCGTTTTACAACGT
CGTGACTGGGAAAACCTGGCGTTACCCAACCTAATCGCCTTGCA
CACATCCCCCTTTCGCCAGCTGGCGTAATAGCGAAGAGGCCCGCAC
CGATCGCCCTTCCCAACAGTTGCCAGCCTGAATGGCGAATGGCGC
TTCGCTTGGTAATAAGCCCGCTTCGGCGGGCTTTTTTTT

SEQUENCE LISTING

<110> XU, Ming-Qun
EVANS, Thomas C.
PRADHAN, Sriharsa
COMB, Donald G.
PAULUS, Henry
SUN, Luo
CHEN, Lixin
GHOSH, Inca
NEW ENGLAND BIOLABS, INC.
BOSTON BIOMEDICAL RESEARCH INSTITUTE

<120> METHOD FOR GENERATING SPLIT, NON-TRANSFERABLE GENES
THAT ARE ABLE TO EXPRESS AN ACTIVE PROTEIN PRODUCT

<130> NEB-163-PCT

<140>
<141>

<150> 60/135,677
<151> 1999-05-24

<160> 134

<170> PatentIn Ver. 2.0

<210> 1
<211> 19
<212> DNA
<213> Escherichia coli

<400> 1
ggacggggaa ctaactatg 19

<210> 2
<211> 20
<212> DNA
<213> Escherichia coli

<400> 2
ccacgatgac gcaccacgcg 20

<210> 3
<211> 30
<212> DNA
<213> Escherichia coli

<400> 3

ggagggggca tatgaatggc gcacagtggg

30

<210> 4

<211> 25

<212> DNA

<213> Escherichia coli

<400> 4

gggggggtcat gataatttct ccaac

25

<210> 5

<211> 28

<212> DNA

<213> Escherichia coli

<400> 5

ccgggtggcg taattatgcc ggtttacg

28

<210> 6

<211> 28

<212> DNA

<213> Escherichia coli

<400> 6

cgtaaaccgg cataattacg ccacccgg

28

<210> 7

<211> 14

<212> PRT

<213> Synechocystis PCC6803

<400> 7

Leu Glu Lys Phe Ala Glu Tyr Cys Phe Asn Lys Ser Thr Gly

1

5

10

<210> 8

<211> 21

<212> PRT

<213> Escherichia coli

<400> 8

Cys Ala Gln Trp Val Val His Ala Leu Arg Ala Gln Gly Val Asn Thr

1

5

10

15

Val Phe Gly Tyr Gly

20

<210> 9

<211> 20

<212> PRT

<213> Escherichia coli

<400> 9

Cys Val Trp Pro Leu Val Pro Pro Gly Ala Ser Asn Ser Glu Met Leu
1 5 10 15

Glu Lys Leu Ser
20

<210> 10

<211> 26

<212> DNA

<213> Escherichia coli

<400> 10

gggggtcatg aatggcgac agtggg

26

<210> 11

<211> 34

<212> DNA

<213> Escherichia coli

<400> 11

gcgcgctcga gttgatttaa cggctgctgt aatg

34

<210> 12

<211> 32

<212> DNA

<213> Escherichia coli

<400> 12

gcgcgaccgg ttgtgactgg cagcaacact gc

32

<210> 13

<211> 31

<212> DNA

<213> Escherichia coli

<400> 13

ggggggctgc agtcataata atttctccaa c

31

<210> 14
<211> 22
<212> DNA
<213> MAIZE

<400> 14
atcagtacac agtcctgcca tc

22

<210> 15
<211> 20
<212> DNA
<213> MAIZE

<400> 15
gagacagccg ccgcaaccat

20

<210> 16
<211> 29
<212> DNA
<213> MAIZE

<400> 16
gggcccataat ggccaccgcc gccgccgcg

29

<210> 17
<211> 29
<212> DNA
<213> MAIZE

<400> 17
gggccctcga ggcttccttc aagaagagc

29

<210> 18
<211> 29
<212> DNA
<213> MAIZE

<400> 18
gggccaccgg tacatcaaag aagagcttg

29

<210> 19
<211> 31
<212> DNA
<213> MAIZE

<400> 19
ggggctgcat tcagtacaca gtcctgcat c

31

<210> 20
<211> 7
<212> PRT
<213> Synechocystis PCC6803

<400> 20
Leu Glu Lys Phe Ala Glu Tyr
1 5

<210> 21
<211> 7
<212> PRT
<213> Synechocystis PCC6803

<400> 21
Cys Phe Asn Lys Ser Thr Gly
1 5

<210> 22
<211> 21
<212> PRT
<213> MAIZE

<400> 22
Cys Lys Gly Ala Asp Ile Leu Val Glu Ser Leu Glu Arg Cys Gly Val
1 5 10 15

Arg Asp Val Phe Ala
20

<210> 23
<211> 21
<212> PRT
<213> MAIZE

<400> 23
Cys Ile Pro Ser Gly Gly Ala Phe Lys Asp Met Ile Leu Asp Gly Asp
1 5 10 15

Gly Arg Thr Val Tyr
20

<210> 24
<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 24

ggatcctaag aaggagatat acccatggaa tcctgcacgt taca

44

<210> 25

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 25

gtcgacgctc tcctgcagtt aggcaggcgt actcattc

38

<210> 26

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 26

gctttgctcc tggcggcttt accttgtggt aaaaccgc

38

<210> 27

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 27

gcggttttac cacaaggtaa agccgccagg agcaaagc

38

<210> 28

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 28

gcccctaaag acacaattat tcgcg

25

<210> 29

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 29

cagcgggcgcc gtcacgagca gagcg

25

<210> 30

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 30

gcgaaccacc actaccaaca atttg

25

<210> 31

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 31

tatctccacg ccaaagggtt tcatt

25

<210> 32

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 32

gaatattgcc tgtcttttgg t

21

<210> 33

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 33

gttaaagcag ttagcagcga t

21

<210> 34

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 34

tgctgaatat tgcctgtctt ttgg

24

<210> 35

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 35

ccgttaaagc agttagcagc gatagc

26

<210> 36

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 36

ggatcctaag aaggagatat acccatggaa tccctgacgt taca

44

<210> 37

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 37

gatatcctgc agttaacctg gagagtgata ctgttgacc

39

<210> 38

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 38

gatatcccat gggacgctat ctggctcagg gcgatg

36

<210> 39

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic,
based on Salmonella typhimurium

<400> 39

gtcgacgctc tcctgcagtt aggcaggcgt actcattc

38

<210> 40

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic from
Synechocystis species PCC6803

<400> 40

tgctgaatat gcgctgtctt ttggtaccga a

31

<210> 41

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic from
Synechocystis species PCC6803

<400> 41

ccgttaaacg ccgcagcagc gatagcgcc

29

<210> 42

<211> 178

<212> PRT

<213> Escherichia coli

<400> 42

Tyr Ala Val Asp Lys Ala Asp Leu Leu Leu Ala Leu Gly Val Arg Phe
1 5 10 15Asp Asp Arg Val Thr Lys Ile Glu Ala Phe Ala Ser Arg Ala Lys Ile
20 25 30Val His Val Asp Ile Asp Pro Ala Glu Ile Gly Lys Asn Lys Gln Pro
35 40 45His Val Ser Ile Cys Ala Asp Val Lys Leu Ala Leu Gln Gly Met Asn
50 55 60Ala Leu Leu Glu Gly Ser Thr Ser Lys Lys Ser Phe Asp Phe Gly Ser
65 70 75 80Trp Asn Asp Glu Leu Asp Gln Gln Lys Arg Glu Phe Pro Leu Gly Tyr
85 90 95Lys Thr Ser Asn Glu Glu Ile Gln Pro Gln Tyr Ala Ile Gln Val Leu
100 105 110

Asp Glu Leu Thr Lys Gly Glu Ala Ile Ile Gly Thr Gly Val Gly Gln
 115 120 125

His Gln Met Trp Ala Ala Gln Tyr Tyr Thr Tyr Lys Arg Pro Arg Gln
 130 135 140

Trp Leu Ser Ser Ala Gly Leu Gly Ala Met Gly Phe Gly Leu Pro Ala
 145 150 155 160

Ala Ala Gly Ala Ser Val Ala Asn Pro Gly Val Thr Val Val Asp Ile
 165 170 175

Asp Gly

<210> 43

<211> 179

<212> PRT

<213> Escherichia coli

<400> 43

Tyr Ala Val Asp Ser Ser Asp Leu Leu Leu Ala Phe Gly Val Arg Phe
 1 5 10 15

Asp Asp Arg Val Thr Gly Lys Leu Glu Ala Phe Ala Ser Arg Ala Lys
 20 25 30

Ile Val His Ile Asp Ile Asp Ser Ala Glu Ile Gly Lys Asn Lys Gln
 35 40 45

Pro His Val Ser Ile Cys Ala Asp Ile Lys Leu Ala Leu Gln Gly Leu
 50 55 60

Asn Ser Ile Leu Glu Ser Lys Glu Gly Lys Leu Lys Leu Asp Phe Ser
 65 70 75 80

Ala Trp Arg Gln Glu Leu Thr Glu Gln Lys Val Lys His Pro Leu Asn
 85 90 95

Phe Lys Thr Phe Gly Asp Ala Ile Pro Pro Gln Tyr Ala Ile Gln Val
 100 105 110

Leu Asp Glu Leu Thr Asn Gly Asn Ala Ile Ile Ser Thr Gly Val Gly
 115 120 125

Gln His Gln Met Trp Ala Ala Gln Tyr Tyr Lys Tyr Arg Lys Pro Arg

130 135 140
 Gln Trp Leu Thr Ser Gly Gly Leu Gly Ala Met Gly Phe Gly Leu Pro
 145 150 155 160
 Ala Ala Ile Gly Ala Ala Val Gly Arg Pro Asp Glu Val Val Val Asp
 165 170 175
 Ile Asp Gly

<210> 44
 <211> 179
 <212> PRT
 <213> Escherichia coli

<400> 44
 Tyr Ala Val Asp Ser Ser Asp Leu Leu Leu Ala Phe Gly Val Arg Phe
 1 5 10 15
 Asp Asp Arg Val Thr Gly Lys Leu Glu Ala Phe Ala Ser Arg Ala Lys
 20 25 30
 Ile Val His Ile Asp Ile Asp Ser Ala Glu Ile Gly Lys Asn Lys Gln
 35 40 45
 Pro His Val Ser Ile Cys Ala Asp Ile Lys Leu Ala Leu Gln Gly Leu
 50 55 60
 Asn Ser Ile Leu Glu Ser Lys Glu Gly Lys Leu Lys Leu Asp Phe Ser
 65 70 75 80
 Ala Trp Arg Gln Glu Leu Thr Val Gln Lys Val Lys Tyr Pro Leu Asn
 85 90 95
 Phe Lys Thr Phe Gly Asp Ala Ile Pro Pro Gln Tyr Ala Ile Gln Val
 100 105 110
 Leu Asp Glu Leu Thr Asn Gly Ser Ala Ile Ile Ser Thr Gly Val Gly
 115 120 125
 Gln His Gln Met Trp Ala Ala Gln Tyr Tyr Lys Tyr Arg Lys Pro Arg
 130 135 140
 Gln Trp Leu Thr Ser Gly Gly Leu Gly Ala Met Gly Phe Gly Leu Pro
 145 150 155 160

Ala Ala Ile Gly Ala Ala Val Gly Arg Pro Asp Glu Val Val Val Asp
 165 170 175

Ile Asp Gly

<210> 45

<211> 180

<212> PRT

<213> Escherichia coli

<400> 45

Met Thr Met His Asn Ala Asp Val Ile Phe Ala Val Gly Val Arg Phe
 1 5 10 15

Asp Asp Arg Thr Thr Asn Asn Leu Ala Lys Tyr Cys Pro Asn Ala Thr
 20 25 30

Val Leu His Ile Asp Ile Asp Pro Thr Ser Ile Ser Lys Thr Val Thr
 35 40 45

Ala Asp Ile Pro Ile Val Gly Asp Ala Arg Gln Val Leu Glu Gln Met
 50 55 60

Leu Glu Leu Leu Ser Gln Glu Ser Ala His Gln Pro Leu Asp Glu Ile
 65 70 75 80

Arg Asp Trp Trp Gln Gln Ile Glu Gln Trp Arg Ala Arg Gln Cys Leu
 85 90 95

Lys Tyr Asp Thr His Ser Glu Lys Ile Lys Pro Gln Ala Val Ile Glu
 100 105 110

Thr Leu Trp Arg Leu Thr Lys Gly Asp Ala Tyr Val Thr Ser Asp Val
 115 120 125

Gly Gln His Gln Met Phe Ala Ala Leu Tyr Tyr Pro Phe Asp Lys Pro
 130 135 140

Arg Arg Trp Ile Asn Ser Gly Gly Leu Gly Thr Met Gly Phe Gly Leu
 145 150 155 160

Pro Ala Ala Leu Gly Val Lys Met Ala Leu Pro Glu Glu Thr Val Val
 165 170 175

Cys Val Thr Gly
 180

<210> 46

<211> 170

<212> PRT

<213> Escherichia coli

<400> 46

Phe Ala Val Gln Glu Cys Asp Leu Leu Ile Ala Val Gly Ala Arg Phe
1 5 10 15

Asp Asp Arg Val Thr Gly Lys Leu Asn Thr Ser Ala Pro His Ala Ser
20 25 30

Val Ile His Met Asp Ile Asp Pro Ala Glu Met Asn Lys Leu Arg Gln
35 40 45

Ala His Val Ala Leu Gln Gly Asp Leu Asn Ala Leu Leu Pro Ala Leu
50 55 60

Gln Gln Pro Leu Asn Gln Cys Asp Trp Gln Gln His Cys Ala Gln Leu
65 70 75 80

Arg Asp Glu His Ser Trp Arg Tyr Asp His Pro Gly Asp Ala Ile Tyr
85 90 95

Ala Pro Leu Leu Leu Lys Gln Leu Ser Asp Arg Lys Pro Ala Asp Cys
100 105 110

Val Val Thr Thr Asp Val Gly Gln His Gln Met Trp Ala Ala Gln His
115 120 125

Ile Ala His Thr Arg Pro Glu Asn Phe Ile Thr Ser Ser Gly Leu Gly
130 135 140

Thr Met Gly Phe Gly Leu Pro Ala Ala Val Gly Ala Gln Val Ala Arg
145 150 155 160

Pro Asn Asp Thr Val Val Cys Ile Ser Gly
165 170

<210> 47

<211> 35

<212> DNA

<213> Escherichia coli

<400> 47

gccttaatta accatgaggg aagcgggtgat cgccg

35

<210> 48

<211> 34

<212> DNA

<213> Escherichia coli

<400> 48

tgcggtcgac ttgcccgact accttggtga tctc

34

<210> 49

<211> 41

<212> DNA

<213> Escherichia coli

<400> 49

cccaagcttg gcgccatgag taaaggagaa gaacttttca c

41

<210> 50

<211> 36

<212> DNA

<213> Escherichia coli

<400> 50

gcgaccgggt tatttgtata gttcatccat gccatg

36

<210> 51

<211> 39

<212> DNA

<213> Escherichia coli

<400> 51

aggggaattcg tcgacaaatt tgctgaatat tgctgtct

39

<210> 52

<211> 38

<212> DNA

<213> Escherichia coli

<400> 52

ggcctcgagt tatttaattg tcccagcgtc aagtaatg

38

<210> 53

<211> 41

<212> DNA

<213> Escherichia coli

<400> 53

agctttggtt aaaccatggt taaagttatc ggtcgtagat c

41

<210> 54

<211> 43

<212> DNA

<213> Escherichia coli

<400> 54

cagcgctcgac ggcgccgtgg gatttggttaa agcagtttagc agc

43

<210> 55

<211> 31

<212> DNA

<213> Escherichia coli

<400> 55

catgccatgg gggaagcggg gatcgccgaa g

31

<210> 56

<211> 39

<212> DNA

<213> Escherichia coli

<400> 56

acgcgagctc ttatttaatt gtcccagcgt caagtaatg

39

<210> 57

<211> 34

<212> DNA

<213> Escherichia coli

<400> 57

cgaattctat ggttaaagtt atcggtcgta gatc

34

<210> 58

<211> 36

<212> DNA

<213> Escherichia coli

<400> 58

agcccgcggt tatttgata gttcatccat gccatg

36

<210> 59

<211> 154

<212> DNA

<213> Nicotiana tabacum

<400> 59

gaatagatct acatacacct tgggtgacac gagtatataa gtcattgtat actgttgaat 60
 aacaagcctt ccattttcta ttttgatttg tagaaaacta gtgtgcttgg gagtcctga 120
 tgattaaata aaccaagatt ttaccttaat taag 154

<210> 60

<211> 151

<212> DNA

<213> Nicotiana tabacum

<400> 60

gacccctggcc tagtctatag gaggttttga aaagaaagga gcaataatca ttttcttgtt 60
 ctatcaagag ggtgctattg ctcttttctt tttttctttt tatttattta ctagtatttt 120
 acttacatag acttttttgt ttacgtattc t 151

<210> 61

<211> 185

<212> DNA

<213> Nicotiana tabacum

<400> 61

catatggcgt ccatgatctc ctctgcccgc gtgaccacgg tcagccgcgc gtccacgggtg 60
 cagtcggccg cggcggcccc gttcggcggc ctcaagtcca tgaccggctt cccgggtcaag 120
 aaggtaaca cggacatcac gtccatcac agcaacggcg gcagggtgaa gtgcatgcga 180
 agagc 185

<210> 62

<211> 6232

<212> DNA

<213> Unknown

<220>

<223> nucleotides 1-2492: E. coli vector pLITMUS28 (New
 England Biolabs, Inc.)

<220>

<223> nucleotides 2493-5993: Nicotiana tabaceum

<220>

<223> Nucleotides 5993-6232: E.coli vector pLITMUS28
 (New England Biolabs, Inc.)

<400> 62

gttaactacg tcagggtggca cttttcgggg aaatgtgcgc ggaacccta tttgtttatt 60
 tttctaaata cattcaaata tgtatccgct catgagacaa taacctgat aaatgcttca 120
 ataattattga aaaaggaaga gtatgagtat tcaacatttc cgtgtcgcgc ttattccctt 180
 ttttgccgca ttttgccctc ctgtttttgc tcaccagaa acgctggtga aagtaaaaga 240
 tgctgaagat cagttgggtg cacgagtggg ttacatcgaa ctggatctca acagcggtaa 300
 gatccttgag agttttcgcc ccgaagaacg ttctccaatg atgagcactt ttaaagttct 360

gctatgtggc gcggtattat cccgtgttga cgccgggcaa gagcaactcg gtcgccgcat 420
acactattct cagaatgact tgggttagta ctcaccagtc acagaaaagc atcttacgga 480
tggcatgaca gtaagagaat tatgcagtgc tgccataacc atgagtata acactgcggc 540
caacttactt ctgacaacga tcggaggacc gaaggagcta accgcttttt tgcacaacat 600
gggggatcat gtaactcgcc ttgatcgttg ggaaccggag ctgaatgaag ccataccaaa 660
cgacgagcgt gacaccacga tgcctgtagc aatggcaaca acgttgcgca aactattaac 720
tggcgaaacta cttactctag cttcccggca acaattaata gactggatgg aggcggataa 780
agttgcagga ccacttctgc gtcgggccct tccggctggc tggtttattg ctgataaatc 840
tggagccggt gagcgtgggt ctcgcggtat cattgcagca ctggggccag atggtaagcc 900
ctcccgtatc gtagttatct acacgacggg gagtcaggca actatggatg aacgaaatag 960
acagatcgct gagatagggt cctcactgat taagcattgg taactgtcag accaagttaa 1020
ctcatatata ctttagattg atttaccctg gttgataatc agaaaagccc caaaaacagg 1080
aagattgtat aagcaaatat ttaaattgta aacgttaata ttttgtaaa attcgcgtta 1140
aatTTTTgtt aaatcagctc atTTTTtaac caataggccg aaatcgga aatcccttat 1200
aaatcaaaag aatagcccg gatagggttg agtggtgttc cagtttgga caagagtcca 1260
ctattaaaga acgtggactc caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc 1320
ccactacgtg aaccatcacc caaatcaagt tttttgggt cgaggtgccg taaagcacta 1380
aatcggaacc ctaaaggag ccccgattt agagcttgac ggggaaagcg aacgtggcga 1440
gaaaggaagg gaagaaagcg aaaggagcgg gcgctagggc gctggcaagt gtagcggta 1500
cgctgcgctg aaccaccaca cccgcgcgc ttaatgcgc gctacagggc gcgtaaaagg 1560
atctagggtg agatcctttt tgataatctc atgacaaaaa tcccttaacg tgagttttcg 1620
ttccactgag cgtagacccc cgtagaaaag atcaaaggat cttcttgaga tccttttttt 1680
ctgcgcgtaa tctgctgctt gcaaacaaaa aaaccacgc taccagcggg ggtttgtttg 1740
ccggatcaag agctaccaac tctttttccg aaggtaactg gcttcagcag agcgagata 1800
ccaaatactg tcttcttagt gtagccgtg ttaggccacc acttcaagaa ctctgtagca 1860
ccgcctacat acctcgctct gctaactctg ttaccagtgg ctgctgccag tggcgataag 1920
tcgtgtctta ccgggttgga ctcaagacga tagttaccgg ataaggcgca gcggtcgggc 1980
tgaacggggg gttcgtgcac acagcccagc ttggagcgaa cgacctacac cgaactgaga 2040
tacctacagc gtgagctatg agaaagcgcc acgcttcccg aaggagagaa ggcggacagg 2100
tatccggtaa gcggcagggg cggaacagga gagcgcacga gggagcttcc agggggaaac 2160
gcctgggtatc tttatagtc tgctgggttt cgccacctct gacttgagcg tcgatttttg 2220
tgatgctcgt cagggggggc gagcctatgg aaaaacgcca gcaacgcggc ctttttacgg 2280
ttcctggcct tttgctggcc ttttgctcac atgtaatgtg agttagctca ctcataggc 2340
accccaggct ttacacttta tgcttccggc tcgtatgttg tgtggaattg tgagcggata 2400
acaatttcac acaggaaaca gctatgacca tgattacgcc aagctacgta atacgactca 2460
ctagtgggca gatcttcgaa tgcacgcgc gcttgacgat atagcaattt tgcttgatt 2520
tatcagtcga agcaggagac aatatacctt gatattctcg atcattctt gattcaaagc 2580
atcgttccat ctcaattgaa aaagcaata acgtttcaag aacaaatcta gttctgcttc 2640
cgtgttgctt ttgtattgtt ttttctttt acccttctt gtgtctgatt ccgcgtaac 2700
tttttaaga gcgttttgat gttttgagag aacagggcc agatttcctt tgtttctat 2760
atctgatcca cgctctttt ctccttgact tgcgggttct tttgcttctt gaattcgatt 2820
ctttattttt ttatttgatc gtagaaaaa gttttgtttt tggtttttat tgatgtttt 2880
attttgacta acattttcat ttgtattcaa atttaaaaga agtaatttgc ttggtataat 2940
ccacggtttt attttatata cattataaag tggtaaaat tctgggaaga accaaaattc 3000
cagattcaat atgggacgat ttaatatatt ttcattcatt cccatccaat caaaaaaggc 3060
ttttttcgaa ttttttgat tgttttctgg attttgatga atcgtaagat aaaaaagcc 3120
ttttttatca attttatcaa ttatttgata attattaata ccaatttttag tatttgatt 3180
actgttggtg togatcttaa cccaggcctc aatatcttct tttgtctaa gagaaaaatg 3240

gataattttc caatcaaaat attttctatc gagatttctt tctatatata gaattattgcc 3300
 ttttcttaga taattattga tatgaagatt gccgagcata tcaaaaaggt tgtgtttgga 3360
 cgtgttggaa ttagaagaaa ttctgagggt cttattttact tgaaagggtg atctagaaat 3420
 aaaagagtca tttttttttt cataattaat cgattttatat gctaaaagat catatctata 3480
 acatttttga aaattatctt tttggtttgc taatgaatag agctcagaat cattttcttt 3540
 tttgtaatga attaatgggt ctttttcata tgaattccat ttgtttaaat ttcgattttg 3600
 agccatacaa ctttgattaa ccttatttcg ccatttttgt ggcattaatc tagaccatct 3660
 aatctgagat aaatcgtacg agaatactca atcatgaata aatgcaagaa aataacctct 3720
 ccttcttttt ctataatgta aacaaaaaag tctatgtaag taaaatacta gtaaataaat 3780
 aaaaagaaaa aaagaaagga gcaatagcac cctcttgata gaacaagaaa atgattattg 3840
 ctcttttctt ttcaaaacct cctatagact aggccaggat cctcgagctt aattaaggta 3900
 aaatcttgggt ttattttaatc atcagggact cccaagcaca ctagttttct acaaatcaaa 3960
 atagaaaata gaaaatggaa ggctttttat tcaacagtat aacatgactt atatactcgt 4020
 gtcaaccaag gtgtatgtag atctattcct gcaggatattc tggatccacg aagcttccca 4080
 tgggaataga tctacatata ccttggttga cagcagatata taagtcatgt tatactgttg 4140
 aataaaaaagc cttccatttt ctattttgat ttgtagaaaa ctagtgtgct tgggagtccc 4200
 tgatgattaa ataaaccaag attttaccgt ttaaaccaccg gtgatcctgg cctagtctat 4260
 aggaggtttt gaaaagaaag gagcaataat cattttcttg ttctatcaag aggggtgctat 4320
 tgctcctttc tttttttctt tttatttatt tactagtatt ttacttacat agactttttt 4380
 gtttacatta tagaaaaaga aggagaggtt attttcttgc atttattcat gattgagtat 4440
 tctcctaggt gtattgataa tgccgtctta accagttttt ccattgattg attctataac 4500
 tctgaagttt cttatgtttt aattcagaat gaaatattcc tagtggtcga aaatagtcct 4560
 ttattttagt cttaaggaaa aaagacgttc tgttatattg aagaacagat cttaatttag 4620
 acaaattaat aacttgggggt tgtgataatt tgtaaaatac atatgcttgt gataagtagg 4680
 ataaatcaaa aaaaatatgt gaatttttct tactaatatt ataaagtac ttttttatag 4740
 tcgaaataaa gtgaattttt ttttgattat taatttttct ttgatttatt tcattattgg 4800
 aaatgtattt atcaatcaat ttgtttgttg attcaagaaa gagttgtgta ttaattctgg 4860
 gaatattaat gatagataaa aatagatcga tgtataatct ttgaatgaat aatttttagaa 4920
 aataatggaa ttccatattt aatcgagtat ttcttctttt taatatttggt aaaaatctttt 4980
 ttggcgattc gaatttttta atattatttg ttttattagg actaatgtct atttctggag 5040
 ttactttctt tttctctttt gtaattcttt ctatttgatt tttgattgta cttgttctat 5100
 cagtcaaatc ctccattttg ctttctatca gtgaagaatt tggccaattt ccagattcaa 5160
 tttgactaaa tgattcgtta attatctgat tactcattag agaatctttt tctttttctg 5220
 tttcattcga ttcatctatt tctttgagtc taaataatac aattggattt acttttgaaa 5280
 gttctttttt catttttttt ataaatagac tacttttgat aagccatttt ttggtttctt 5340
 ttgaaattct tcgaaataat tttatttttc ctttgaaaac ttttagagtt ataaaatatt 5400
 tctttttgaa ttttccaatt tttttttcga gttccttaaa aatggggtca aaaaaagaag 5460
 ggcgttttct gggagaacca aagggaagtt cagcttccat tccccaaact gttaaaaaac 5520
 aaaaatcatc tttttgtttt ttctttttca ttagctctcc acgggaggag tacagttagg 5580
 atatatgcca aggtttcaga caaaaaggaa ataataattt gatctgaatg ccatctttca 5640
 accaattttt tggaaattct gtttctgata attgaacacc attataagta catttaatat 5700
 gcatttctct attccattcc tgcaaatctt cagaccattc aggaagttgc aagactaaca 5760
 tacgcccgag atttttggct attatcaatg aaggtaatac aatatatttt cgaagaattg 5820
 attgagttat taacatgtaa cctcttatta tttgcgcaa aggaatggta tcccaggctt 5880
 ctgctatctc tatcgtgtct ttttctttc ttttgttctc cctttttttg tcttttctt 5940
 ttttctcttc tctttttgtt tgttcttctc tagactctag aatcttgaat tcggtaccct 6000
 ctagtcaagg ccttaagtga gtcgtattac ggactggccg tcgttttaca acgtcgtgac 6060
 tgggaaaacc ctggcggtac ccaacttaat cgccttgacg cacatcccc tttcgccagc 6120

tggcgtaata gcgaagaggc ccgcaccgat cgccttccc aacagttgcg cagcctgaat 6180
ggcgaatggc gcttcgcttg gtaataaagc ccgcttcggc gggctttttt tt 6232

<210> 63

<211> 6477

<212> DNA

<213> Unknown

<220>

<223> Nucleotides 1-2482: E. coli vector pLITMUS28 (New
England Biolabs, Inc.)

<220>

<223> Nucleotides 2493-6242: Nicotiana tabaceum

<220>

<223> Nucleotides 6243-6477: E. coli vector pLITMUS28
(New England Biolabs, Inc.)

<400> 63

gttaactacg tcaggtggca cttttcgggg aaatgtgcmc ggaacccta tttgtttatt 60
tttctaaata cattcaaata tgtatccgct catgagacaa taacctgat aaatgcttca 120
ataatattga aaaaggaaga gtatgagtat tcaacatttc cgtgtcgccc ttattccctt 180
ttttgcgga ttttgccctc ctgtttttgc tcaccagaa acgctgggtga aagtaaaaga 240
tgctgaagat cagttgggtg cagcagtggt ttacatcgaa ctggatctca acagcggtaa 300
gatccttgag agttttcgcc ccgaagaacg ttctccaatg atgagcactt ttaaagttct 360
gctatgtggc gcggtattat cccgtgttga cgcggggcaa gagcaactcg gtcgcccgat 420
acactattct cagaatgact tggttgagta ctcaccagtc acagaaaagc atcttacgga 480
tgccatgaca gtaagagaat tatgcagtgc tgccataacc atgagtata aactgcccgc 540
caacttactt ctgacaacga tcggaggacc gaaggagcta accgcttttt tgcacaacat 600
gggggatcat gtaactcgcc ttgatcggtg ggaaccggag ctgaatgaag ccataccaaa 660
cgacgagcgt gacaccacga tgccctgtagc aatggcaaca acgttgcgca aactattaac 720
tgccgaacta ctactctag cttcccggca acaattaata gactggatgg aggcggataa 780
agttgcagga ccacttctgc gtcgcccct tccggctggc tggtttattg ctgataaatc 840
tgagccgggt gagcgtgggt ctgcgggtat cattgcagca ctggggccag atggttaagc 900
ctcccgtatc gtagttatct acacgacggg gagtcaggca actatggatg aacgaaatag 960
acagatcgct gagatagggt cctcactgat taagcattgg taactgtcag accaagtta 1020
ctcatatata ctttagattg atttaccocg gttgataatc agaaaagccc caaaaacagg 1080
aagattgtat aagcaaata ttaaattgta aacgttaata tttgttaaa attcgcgtta 1140
aatttttgtt aaatcagctc attttttaac caataggccg aaatcggcaa aatcccttat 1200
aaatcaaaag aatagcccga gatagggttg agtgtgttc cagtttggaa caagagtcca 1260
ctattaaaga acgtggactc caacgtcaaa gggcgaaaaa ccgtctatca gggcgatggc 1320
ccactacgtg aaccatcacc caaatcaagt tttttgggtg cgaggtgccg taaagcacta 1380
aatcggaacc ctaaaggag ccccgattt agagcttgac ggggaaagcg aacgtggcga 1440
gaaaggaagg gaagaaagcg aaaggagcgg gcgctaggcg gctggcaagt gtagcggta 1500
cgctgcgcgt aaccaccaca cccgcgcgc ttaatgcgc gctacaggcg gcgtaaaagg 1560
atctagggtg agatcctttt tgataatctc atgacaaaaa tcccttaacg tgagttttcg 1620
ttccactgag cgtcagaccc cgtagaaaag atcaaaggat cttcttgaga tccttttttt 1680

ctgcgcgtaa tctgctgctt gcaaacaaaa aaaccaccgc taccagcggt ggtttgtttg 1740
 ccggatcaag agctaccaac tctttttccg aaggttaactg gcttcagcag agcgcagata 1800
 ccaaatactg ttctttctagt gtagccgtag ttaggccacc acttcaagaa ctctgtagca 1860
 ccgcctacat acctcgctct gctaatacctg ttaccagtgg ctgctgccag tggcgataag 1920
 tcgtgtctta ccgggttgga ctcaagacga tagttaccgg ataaggcgca gcggtcgggc 1980
 tgaacggggg gttcgtgcac acagcccagc ttggagcgaa cgacctacac cgaactgaga 2040
 tacctacagc gtgagctatg agaaagcgcc acgcttcccg aaggagaaaa ggcgacagg 2100
 tatccggtaa gcggcagggt cggaacagga gagcgcacga gggagcttcc agggggaaac 2160
 gcctggatc tttatagtcc tgtcgggttt cgccacctct gacttgagcg tcgatttttg 2220
 tgatgctcgt caggggggcg gagcctatgg aaaaacgcca gcaacgcggc ctttttacgg 2280
 ttcttgccct tttgctggcc ttttgctcac atgtaatgtg agttagctca ctcattaggc 2340
 accccaggct ttacacttta tgcttccggc tcgtatgttg tgtggaattg tgagcggata 2400
 acaatttcac acaggaataa gctatgacca tgattacgcc aagctacgta atacgactca 2460
 ctagtgggca gatcttcgaa tgcacgcgc gcaattcacc gccgtatggc tgaccggcga 2520
 ttactagcga ttccggcttc atgcaggcga gttgcagcct gcaatccgaa ctgaggacgg 2580
 gtttttgggg ttagctcacc ctccgggat cgcgaccctt tgtcccgcc attgtagcac 2640
 gtgtgtcgcc cagggcataa ggggcatgat gacttgacgt catcctcacc ttctccggc 2700
 ttatcaccgg cagtctgttc agggttccaa actcaacgat ggcaactaaa cacgagggtt 2760
 gcgctcgttg cgggacttaa cccaacacct tacggcacga gctgacgaca gccatgcacc 2820
 acctgtgtcc gcgttcccgaggacccct ctctttcaag aggattcgcg gcatgtcaag 2880
 ccttggttaag gttcttcgct ttgcatcgaa ttaaaccaca tgctccaccg cttgtcggg 2940
 ccccgctcaa ttcttttgag ttctattctt gcgaacgtac tcccaggcg ggatacttaa 3000
 cgcgtttagct acagcactgc acgggtcgat acgcacagcg cctagtatcc atcgtttacg 3060
 gctaggacta ctggggatc taatcccatt cgctccctta gctttcgtct ctcagtgtca 3120
 gtgtcgggcc agcagagtgc ttccgcggtt ggtgttcttt ccgatctcta cgcatctcac 3180
 cgctccaccg gaaattccct ctgcccctac cgtactccag cttggtagt tccaccgct 3240
 gtccagggtt gagccctggg atttgacggc ggacttaaaa agccacctac agacgcttta 3300
 cgcccaatca ttccggataa cgcttgatc ctctgtatta ccgcggtgc tggcacagag 3360
 ttagccgatg cttattcccc agataccgctc attgcttctt ctccgggaaa agaagttcac 3420
 gaccggtggg ctttctacct ccacgcggca ttgctccgct agctttcgcc cattgcggaa 3480
 aattccccac tgctgcctcc cgtaggagtc tgggcccgtgt ctcagtccca gtgtggctga 3540
 tcatcctctc ggaccagcta ctgatcatcg ccttggttaag ctattgcctc accaactagc 3600
 taatcagacg cgagccccct ctccggcgga ttctctctt tgctcctcag cctacgggg 3660
 attagcagcc gtttccagct gttgttcccc tcccaagggc aggttcttac gcgttactca 3720
 ccgctccgcc actggaaaca ccacttcccg tccgacttgc atgtgttaag catgccgcca 3780
 gcgttcatcc tgagccagga tcgaactctc catgagatc atagtgtcat tacttatagc 3840
 ttcttgttc gtagacaaag cggattcgga attgtcttct attccaaggc ataacttgta 3900
 tccatgcgct tcatattcgc cgggagttcg ctcccagaaa tatagccatc cctgccccct 3960
 cagtcatac ccacgagcct cttatccatt ctcatgaac gacggcgggg gagcaaatcc 4020
 aactagaaaa actcacattg ggcttaggga taatcaggct cgaactgatg acttccacca 4080
 cgtcaagggt acactctacc gctgagttat atcccttccc cgccccatcg agaaatagaa 4140
 ctgactaatc ctaagtcaaa ggcgtacgag aatactcaat catgaataaa tgcaagaaaa 4200
 taacctctcc ttctttttct ataagtataa caaaaaagtc tatgtaagta aaatactagt 4260
 aaataaataa aaagaaaaaa agaaaggagc aatagacccc tcttgataga acaagaaaat 4320
 gattattgct ctttctttt caaaacctcc tatagactag gccaggatcc tcgagcttaa 4380
 ttaaggtaaa atcttggttt atttaatcat cagggactcc caagcacact agttttctac 4440
 aaatcaaaat agaaaataga aaatggaagg ctttttatte aacagtataa catgacttat 4500
 atactcgtgt caaccaagggt gtatgtagat ctattcctgc aggatatctg gatccacgaa 4560

```

gcttcccatg ggaatagatc tacatacacc ttggttgaca cgagtatata agtcattgta 4620
tactgttgaa taaaaagcct tccattttct attttgattt gtagaaaact agtggtgcttg 4680
ggagtccttg atgattaaat aaaccaagat tttaccgttt aaacaccggg gatcctggcc 4740
tagtctatag gaggttttga aaagaaagga gcaataatca ttttcttggt ctatcaagag 4800
ggtgctattg ctccctttct tttttctttt tatttattta ctagtatttt acttacatag 4860
acttttttgt ttacattata gaaaaagaag gagagggttat tttcttgcat ttattcatga 4920
ttgagtattc tcctaggggc gagaaactca acgccactat tcttgaacaa cttggagccg 4980
ggccttcttt tcgcactatt acggatatga aaataatggg caaaatcgga ttcaattgtc 5040
aactgcccct atcggaataa ggattgacta ccgattccga aggaactgga gttacatctc 5100
ttttccattc aagagttctt atgcgtttcc acgccccttt gagaccccga aaaatggaca 5160
aattcctttt cttaggaaca catacaagat tcgtcactac aaaaaggata atggtaaccc 5220
taccattaac tacttcattt atgaatttca tagtaataga aatacatgtc ctaccgagac 5280
agaatttggg acttgctatc ctcttgcta gcaggcaaag atttacctcc gtggaaagga 5340
tgattcattc ggatcgacat gagagtccaa ctacattgcc agaatccatg ttgtatattt 5400
gaaagagggt gacctccttg cttctctcat ggtacactcc tcttcccgcc gagccccctt 5460
tctcctcggt ccacagagac aaaaagttag actggtgcca acaattcatc agactacta 5520
agtcgggata actaactaat actaatctaa tataatagtc taatatatct aatataatag 5580
aaaatactaa tataatagaa aagaactgtc ttttctgtat actttccccg gttccgttgc 5640
taccgcgggc tttacgcaat cgatcggatt agatagatat ccttcaaca taggtcatcg 5700
aaaggatctc ggagaccac caaagtacga aagccaggat ctttcagaaa acggattcct 5760
attcaaagag tgcataaccg catggataag ctcacactaa cccgtcaatt tgggatccaa 5820
attcgagatt ttccttgga ggtatcgga aggatttga atggaataat atcgattcat 5880
acagaagaaa aggttctcta ttgattcaaa cactgtacct aacctatggg atagggatcg 5940
aggaagggga aaaaccgaag atttcacatg gtacttttat caatctgatt tatttcgtac 6000
ctttcgttca atgagaaaat gggtaaaatt ctacaggatc aaacctatgg gacttaagga 6060
atgatataaa aaaaagagag ggaatatatt catattaaat aaatatgaag tagaagaacc 6120
cagattccaa atgaacaaat tcaaacttga aaaggatctt ccttattctt gaagaatgag 6180
gggcaaaggg attgatcaag aaagatcttt tggtcttctt atatataaga tcgtgatggg 6240
acctctagt caaggcctta agtgagtcgt attacggact ggccgtcgtt ttacaacgtc 6300
gtgactggga aaacctggc gttacccaac ttaatcgct tgcagacat cccctttcg 6360
ccagctggcg taatagcgaa gagggccgca ccgatcgccc tcccaacag ttgcgcagcc 6420
tgaatggcga atggcgcttc gcttggtaat aaagcccgtc tcggcgggct tttttt 6477

```

<210> 64

<211> 31

<212> DNA

<213> Nicotiana tabacum

<400> 64

aactgcagga atagatctac atacaccttg g

31

<210> 65

<211> 42

<212> DNA

<213> Nicotiana tabacum

<400> 65

ccgctcgagc ttaattaagg taaaatcttg gtttatttaa tc

42

<210> 66

<211> 33

<212> DNA

<213> Nicotiana tabacum

<400> 66

gcgaccggtg atcctggcct agtctatagg agg

33

<210> 67

<211> 34

<212> DNA

<213> Nicotiana tabacum

<400> 67

aggcctagga gaataactcaa tcatgaataa atgc

34

<210> 68

<211> 34

<212> DNA

<213> Nicotiana tabacum

<400> 68

ttggcgcgct tgacgatata gcaattttgc ttgg

34

<210> 69

<211> 34

<212> DNA

<213> Nicotiana tabacum

<400> 69

ttgcgtacga tttatctcag attagatggt ctag

34

<210> 70

<211> 35

<212> DNA

<213> Nicotiana tabacum

<400> 70

ttgcctaggc gtattgataa tgccgtctta accag

35

<210> 71

<211> 34

<212> DNA

<213> Nicotiana tabacum

<400> 71

aggggtaccg aattcaagat tctagagtct agag

34

<210> 72

<211> 34

<212> DNA

<213> Nicotiana tabacum

<400> 72

ttggcgcgca attcacgcgc gtatggctga ccgg

34

<210> 73

<211> 34

<212> DNA

<213> Nicotiana tabacum

<400> 73

ttgcgtacgc ctttgactta ggattagtca gtcc

34

<210> 74

<211> 34

<212> DNA

<213> Nicotiana tabacum

<400> 74

ttgcctaggg tcgagaaact caacgccact attc

34

<210> 75

<211> 35

<212> DNA

<213> Nicotiana tabacum

<400> 75

aggggtacca tcacgatctt atatataaga agaac

35

<210> 76

<211> 250

<212> DNA

<213> Nicotiana tabacum

<400> 76

gaattgtgag cgctcacaat tctaggatgt taattgcgcc gacatcataa cggttctggc 60
aaatattctg aaatgagctg ttgacaatta atcatcggtc cgtataatgt gtggaattgt 120
gagcggataa caatttcaca caggaaacag accatgggtga attctagagc tcgaggatcc 180
gcggtaccgg ggcattgcatt cgaagcttcc ttaagcggcc gtcgaccgat gcccttgaga 240
gccttcaacc 250

<210> 77

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 77

Cys Leu Asn Ile Gln

1 5

<210> 78

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 78

Val Phe Lys His Ala

1 5

<210> 79

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 79

Leu Phe Lys Gln Pro

1 5

<210> 80

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 80

Cys Leu Asn Ser Asp

1

5

<210> 81

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 81

Cys Leu Asn Ile Ser

1

5

<210> 82

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 82

Cys Leu Asn Thr Asp

1

5

<210> 83

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 83

Cys Leu Asn Asn Arg

1

5

<210> 84

<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 84
Cys Leu Asn Ser Cys
1 5

<210> 85
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 85
Cys Leu Asn Ser Asp
1 5

<210> 86
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 86
Cys Leu Asn Thr Leu
1 5

<210> 87
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: based on the

ends of the Tn7 transposon

<400> 87

Val Phe Lys Gln Pro

1

5

<210> 88

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 88

Cys Leu Asn Ser Met

1

5

<210> 89

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 89

Cys Leu Asn Asn Tyr

1

5

<210> 90

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 90

Cys Leu Asn Met Ala

1

5

<210> 91
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 91
Val Phe Lys His Lys
1 5

<210> 92
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 92
Cys Leu Asn Thr Lys
1 5

<210> 93
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 93
Cys Leu Asn Lys Asp
1 5

<210> 94
<211> 5
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 94

Met Phe Lys Gln Ile

1 5

<210> 95

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 95

Cys Leu Asn Ile Ile

1 5

<210> 96

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 96

Leu Phe Lys His Glu

1 5

<210> 97

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 97

Val Phe Lys His Phe

1

5

<210> 98

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 98

Cys Leu Asn Ser Val

1

5

<210> 99

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 99

Val Phe Lys Gln Ile

1

5

<210> 100

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 100

Met Phe Lys Gln Ala

1

5

<210> 101

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 101

Leu Phe Lys His His
1 5

<210> 102

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 102

Leu Phe Lys His Gln
1 5

<210> 103

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 103

Met Phe Lys His Val
1 5

<210> 104

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 104

Val Phe Lys Gln Lys

1 5

<210> 105

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 105

Leu Phe Lys Gln Gln

1 5

<210> 106

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 106

Leu Phe Lys His Ser

1 5

<210> 107

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 107

Cys Leu Asn Thr Gly

1 5

<210> 108

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 108

Cys Leu Asn Ser Arg

1

5

<210> 109

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 109

Val Phe Lys His Leu

1

5

<210> 110

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 110

Cys Leu Asn Asn Ile

1

5

<210> 111

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the

ends of the Tn7 transposon

<400> 111

Leu Phe Lys His Gln

1

5

<210> 112

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 112

Cys Leu Asn Lys His

1

5

<210> 113

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 113

Met Phe Lys Gln Tyr

1

5

<210> 114

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 114

Cys Leu Asn Lys Gln

1

5

<210> 115

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 115

Cys Leu Asn Met Ser

1 5

<210> 116

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 116

Leu Cys Leu Asn Ile Leu Ala

1 5

<210> 117

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 117

Asn Cys Leu Asn Ile Asn Ala

1 5

<210> 118

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 118

Leu Met Phe Lys His Leu Ser

1

5

<210> 119

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 119

Thr Leu Phe Lys His Thr Arg

1

5

<210> 120

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 120

Lys Val Phe Lys Gln Lys Glu

1

5

<210> 121

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 121

His Leu Val Phe Lys His Leu

1

5

<210> 122

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 122

Leu Cys Leu Asn Thr Leu Leu

1

5

<210> 123

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 123

Leu Cys Leu Asn Asn Leu Val

1

5

<210> 124

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 124

Glu Val Phe Lys His Glu Gly

1

5

<210> 125

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 125

Lys Val Phe Lys Gln Lys Gly

1

5

<210> 126

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 126

Thr Cys Leu Asn Thr Thr Ile

1

5

<210> 127

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 127

Met Cys Leu Asn Asn Met Asn

1

5

<210> 128

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 128

Leu Leu Phe Lys Gln Leu Arg

1

5

<210> 129

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 129

Arg Cys Leu Asn Asn Arg Leu

1

5

<210> 130

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 130

Met Val Phe Lys Gln Met Ala

1

5

<210> 131

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 131

Ala Met Phe Lys Gln Ala Thr

1

5

<210> 132

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 132

Leu Val Phe Lys His Leu Asp

1

5

<210> 133

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 133

Lys Met Phe Lys Gln Lys Thr

1

5

<210> 134

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: based on the
ends of the Tn7 transposon

<400> 134

Tyr Cys Leu Asn Asn Tyr Phe

1

5

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/14122

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12N 15/10, 15/62, 15/64; C12P 21/02

US CL : 435/69.1, 91.42; 536/23.4

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 91.42; 536/23.4

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,834,247 A (COMB et al) 10 November 1998 (10.11.98), columns 2-3, 8-19.	1-6, 12-13, 16, 21
A	CHONG et al. "Single-column purification of free recombinant proteins using a self-cleavable affinity tag derived from a protein splicing element," Gene. 1997. Vol. 192, pages 271-281, see entire document.	1-32
A	XU. M-Q. "The IMPACT of Protein Splicing Research," The NEB Transcript. January 1997. Vol. 8, No. 2, pages 1-5, see entire document.	1-32

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"g" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

07 AUGUST 2000

Date of mailing of the international search report

07 SEP 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. : (703) 305-3230

Authorized officer

TERRY A. MCKELVEY

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/14122

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SOUTHWORTH et al. "Control of protein splicing by intein fragment reassembly," The EMBO Journal. 1998. Vol. 17, No. 4, pages 918-926, see entire document.	1-32

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/14122

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

West (US and European databases), Dialog OneSearch (biotech databases)

search terms: intein, inteins, extein, exteins, als, acetolactate, epsps, enolpyruvyl, separat?, split?, protein splic?, affinity, compartment? complement?, linker?, loop?

THIS PAGE BLANK (125710)

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)